

GOVERNMENT OF INDIA

PATENT OFFICE BRANCH

~~81, Wallajah Road,~~

~~MADRAS-600 002.~~

GOVERNMENT OF INDIA
PATENT OFFICE BRANCH

WING 'C' - (C-4 A)

III FLOOR, RAJANI BHAVAN
BESANT NAGAR, CHENNAI-600 030.

ORIGINAL

9723

Patent Application No:

708/MAS/97

Date - - 19

03-4-97

Received documents purporting to be an Application for a Patent by
BOARD OF REGENTS, UNIVERSITY OF TEXAS SYSTEM.
relating to Calixpyrrole, Calixpyridinopyrrole And
Calixpyridine macrocycles & 9

together with Provisional Specification and a fee of Rs. 1000 (Rupees One Thousand
and Five Hundred only.) 1,500/-

The Application has been numbered and dated as of 708/MAS/97 dated 03-4-97

To
M/S. De Bevoise & De Bevoise,
B.S. B. Ed., Chen. 600 028.

M. S. Venkataraman

for Controller of Patents
(M. S. VENKATARAMAN)

2nd Asst. Controller of Patents & Designs

(See Notice on the reverse)

Sidel

NOTICE

- (1) An Application accompanied by Provisional Specification will not be examined under Section 12 of the Patents Act, 1970 (39 of 1970) until the Complete Specification accompanied by the prescribed fee of Rs. 200/- is filed.
- (2) The normal period allowable for filing the Complete Specification is **Twelve months** from the date of filing the application. An extension upto three months beyond the said twelve months may be obtained on making a request on Form No. 5 accompanied by a fee of Rs. 100/- per month.
- (3) Where a Complete Specification has been filed in respect of an application, the examination report will be forwarded to the applicant (or his agents) in due course.
- (4) Where an application has been filed with a Provisional Specification, if the Complete Specification is not filed within the time allowed under the Act, the application shall be deemed to have been abandoned under Section 9(1) of the Act.
- 5) If the application is accompanied by a Provisional Specification, a declaration as to inventorship on Form 6 should be filed along with the Complete Specification.

Side 2

DESCRIPTION

CALIX[n]PYRROLES, CALIX[m]PYRIDINO[n]PYRROLES AND CALIX[m]PYRIDINES

BACKGROUND OF THE INVENTION

The present invention relates to ion and neutral molecule binding, separation, or detection, and predicted on the use of calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine macrocycles.

Anions play essential roles in biological processes; indeed, it is believed that they participate in 70% of all enzymatic reactions. There is, therefore, intense effort being devoted to the problem of anion complexation and recognition. In the molecular recognition arena, a number of research groups have followed Nature's lead and have designed and synthesized receptors that use hydrogen bonds alone, or in concert with electrostatic interactions, to coordinate to anions. Nonetheless, there remains at present a critical need for additional anion complexing agents that are either easy to make or inherently selective in their substrate binding properties.

Additionally, the separation of anionic mixtures using chromatographic techniques involving anion binding interactions is a little studied area of chemistry. Current techniques for purification of anions such as oligonucleotide fragments, polyphosphate-containing molecules such as ATP, ADP and AMP, carboxylates, or *N*-protected amino acids involve derivatization prior to separation, leading to decreased yields and cumbersome methodology, or involve a salt-separation step following chromatography. This is a very important area to both scientists and clinicians as both mono- and di-nucleotides, natural and synthetic oligonucleotides play critical roles in modern biotechnology as well as medicine. Oligonucleotides are used, for instance, as hybridization probes in blot analyses, primers for PCR amplification, and for site-specific mutagenesis. Furthermore, in some areas, oligonucleotide-derived products are currently being used as probes for the detection of genetic diseases and for proviral HIV, the causative agent of Acquired Immunodeficiency Syndrome (AIDS). Oligonucleotides are also being considered as potential chemotherapeutic agents, both directly, i.e., in gene therapy, and in an antisense fashion.

The above mentioned applications require oligonucleotide materials of impeccable purity (often greater than > 99.99%). Such purity, however, is not so easy to obtain using existing technology. Presently, gene products and other oligonucleotide-type materials are purified using polyacrylamide gel electrophoresis (PAGE). This approach suffers from the drawback of being time-consuming and low yielding. It also requires toxic materials and is painfully manpower-demanding and low-yielding.

Liquid chromatographic techniques, in particular, high performance liquid chromatography (HPLC) and High Performance Affinity Chromatography (HPAC) employing

specialty silica gels, are currently used to separate biological molecules. Indeed, silica gel phases with bonded groups, such as linear hydrocarbons, amino groups, cyano-groups, carboxylic amides and amino acids are all known. Unfortunately, few if any of these phases are efficacious for the efficient, high-yielding separation of nucleotides and oligonucleotides.

5 Those that work best for this purpose are ion exchange columns which, on a limited basis, can sometimes separate oligonucleotides containing 40 or fewer residues. However, this technique still suffers from several limitations, not least of which being that it requires severe conditions, such as elution at pH 2.7, for routine operation. The use of high concentration buffers (greater than 1M), and gradients which often include formic acid or formamide, also
10 limits the half lives of ion exchange columns. Reverse phase columns are designed for pressures up to 5000 psi. However, this separative technique is based on hydrophobic interaction and, because there is not a specific interaction for phosphorylated species, it is not surprising that effective separation is not generally achieved.

 A type of stationary phase has been described in which purine and pyrimidine bases are
15 bonded to silica gel. With this support, base-pairing interactions between the nucleic acid base pairs and the modified silica gel were expected to improve resolution in nucleic acid separation. Although such stationary phases were used to separate nucleic acid free-bases, purine alkaloids, nucleosides, and mono- and oligonucleotides, this approach unfortunately has demonstrated the least success in the case of oligonucleotide separation, which is the most important area. Thus,
20 prior to the present invention, there remained a critical need for improved solid supports that would effectively separate nucleotides and oligonucleotides.

 Additionally sapphyrin (an expanded porphyrin known to bind anions) has been attached to a functionalized silica gel. The separatory properties of this material were not sufficient to separate oligonucleotides successfully due to broad peaks for (3- to 9-mer)
25 oligonucleotide mixtures. Furthermore, synthetic challenges are associated with preparing the requisite functionalized sapphyrins since they require over 20 synthetic steps to produce.

 Current technology for dialysis in medical applications relies on membranes, such as microfiltering cellophane, to filter anions such as chloride anion or phosphate-containing molecular anions from the blood stream. Aluminum hydroxide or calcium carbonate cocktails
30 must be consumed by the dialysis patient in order to bind the anionic species. A major drawback of this technology is that aluminum build up in cellular membranes to toxic levels over time causing ailments including dementia and death. Calcium carbonate offers a less toxic substitute, however, it is less efficient and is associated with hypercalcemia.

 Water- soluble anion binding agents are desired as drug delivery agents. For example,
35 many anti-viral drugs only show activity when phosphorylated. However the phosphorylated drug derivatives are too polar to pass through cell wall membranes. A water soluble anion chelating agent may be able to encapsulate the negative charge and so allow the drug to pass through cell walls.

The synthesis of new molecular devices designed to sense and report the presence of a particular substrate is an area of analytical chemistry which is attracting attention. The detection of anionic species is a particular challenge, as anions are difficult to bind and are generally larger than cations leading to a smaller charge-to-radius ratio.

5 Molecular recognition of neutral compounds presents a challenge in the area of supramolecular chemistry. Binding of substrates such as short-chain alcohols and simple monoamides is particularly difficult, because these molecules have few functionalized sites available for hydrogen bonding, and they lack the large hydrocarbon surfaces necessary to participate in efficient hydrophobic or π - π stacking interactions. Association constants for
10 neutral substrate-synthetic receptor complexes are thus generally modest, even though the architectural complexity of the receptors is often high.

One aspect of the present invention involves calixpyrroles. This is a subset of a class of macrocycles that was previously termed porphyrinogens. Porphyrinogens are non-conjugated macrocyclic species composed of four pyrrole rings linked in the α -position *via* sp^3 hybridized
15 carbon atoms. Porphyrinogens that carry *meso*-hydrogen atoms are prone to oxidation to the corresponding porphyrins. Fully *meso*-substituted porphyrinogens are generally stable crystalline materials. The first such macrocycle, an octamethyl derivative, was obtained over a century ago by Baeyer (*Ber. Dtsch. Chem. Ges.* 1886, 19, 2184), as the result of an acid-catalyzed condensation between acetone and pyrrole. Subsequently, cyclohexanone was
20 reacted with pyrrole in a 1:1 ratio in the presence of acid to produce tetraspirocyclohexylporphyrinogen. Metal cation binding of the macrocycle is known. A further prior art method uses pyrrole, a C4-C6 saturated alicyclic ketone and an acid containing vinyl groups or triple bonds to form a polymerized resin. In this case, the resulting products are undefined since it appears to be unknown where the modifying group is attached to the
25 product.

While the term porphyrinogen is now widely accepted in the literature, the present inventors believe the fully *meso*-substituted systems with non-hydrogen atoms are misnamed. They are not *bona fide* precursors of the porphyrins and might, therefore, be better considered as being *calixpyrroles*. Such a renaming has precedent in the chemistry of other heterocyclic
30 ring systems. Interestingly, while related functionalized calixarenes, a class of molecules containing benzene as opposed to pyrrole, subunits, have been shown to be capable of binding anions, unmodified calixarene frameworks show no affinity for anionic guests demonstrating that the binding properties are not an inherent part of the calixarene ring structure.

While a few other fully *meso*-substituted porphyrinogens are known, no one prior to
35 the present inventors recognized their properties and consequent uses as described herein.

SUMMARY OF THE INVENTION

The present invention results from the inventors' discovery that calix[4]pyrrole macrocycles bind anions in the solid state. That discovery led the present inventors to synthesize novel calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridine macrocycles. to develop new methods of making said macrocycles, and to demonstrate methods of using said macrocycles as presented herein.

"Calix[n]pyrrole," as used herein, means a macrocycle having "n" or "m+n" pyrrole rings linked in the α positions *via* sp^3 hybridized *meso* carbon atoms that are not bound to hydrogen (protium), deuterium or tritium atoms. This is a subset of a class of molecules that was previously termed porphyrinogens. "Calix[m]pyridino[n]pyrrole," as used herein, means a macrocycle, derived from a calix[m+n]pyrrole, having "m" pyridine rings and "n" pyrrole rings linked in the α positions *via* sp^3 hybridized *meso* carbon atoms that are not bound to hydrogen (protium), deuterium or tritium atoms. "Calix[m]pyridine," as used herein, means a macrocycle, derived from a calix[n]pyrrole, having "m" pyridine rings linked in the α positions *via* sp^3 hybridized *meso* carbon atoms that are not bound to hydrogen (protium), deuterium or tritium atoms.

A "calix[n]pyrrole", "calix[m]pyridino[n]pyrrole" or "calix[m]pyridine" is different from a porphyrinogen since a porphyrinogen has one or more sp^3 hybridized *meso* carbon atoms bound to a hydrogen (protium), deuterium or tritium atom. A "calix[n]pyrrole", "calix[m]pyridino[n]pyrrole" or "calix[m]pyridine" is different from a porphomethene since a porphomethene contains three sp^3 hybridized *meso* carbons and one sp^2 hybridized *meso* carbon. A "calix[n]pyrrole", "calix[m]pyridino[n]pyrrole" or "calix[m]pyridine" is different from a phlorin since a phlorin contains one sp^3 hybridized *meso* carbon and three sp^2 hybridized *meso* carbons. A "calix[n]pyrrole", "calix[m]pyridino[n]pyrrole" or "calix[m]pyridine" is different from a porphyrin since a porphyrin contains four sp^2 hybridized *meso* carbons and is aromatic. A "calix[n]pyrrole", "calix[m]pyridino[n]pyrrole" or "calix[m]pyridine" is different from an expanded porphyrin since an expanded porphyrin contains at least one sp^2 hybridized *meso* carbon.

Macrocycles of the present invention have unexpected properties that make them particularly useful. Calix[n]pyrroles and calix[m]pyridino[n]pyrroles bind anions and neutral molecular species in solution and in the solid state in such an effective and selective way that the anions or neutral molecular species can be separated from other anions and neutral molecular species. Calix[m]pyridino[n]pyrroles and calix[m]pyridines can also be used to bind and separate cations. Further, the affinity a macrocycle has for a particular species can be "tuned" by strategic choice of electron-donating or electron-withdrawing peripheral substituents for synthesis of the macrocycle. Applications of these properties for removal of biological ions or neutral molecule species for medical uses, or removal of undesirable ions or neutral molecule


species from environmental sources provide only a few of the practical and important uses for the present molecules.


5 A calix[n]pyrrole macrocycle noncovalently-complexed to a molecular or anionic species is an embodiment of the present invention. The term "n" refers to the number of pyrrole rings in the macrocycle and, for molecules of the present invention, is 4, 5, 6, 7, or 8.
10 "Noncovalently-complexed to a molecular or anionic species," as used herein, means that bound molecules are held to the core of a macrocycle by noncovalent binding to one or more pyrrole N-H groups. "Noncovalent binding" includes intermolecular interactions such as hydrogen bonds, dipole-dipole interactions, dipole-induced dipole interactions, ion-dipole interactions, ion pairing, Van der Waals interactions, London dispersion forces, π - π stacking interactions, edge-to-face π -interactions, cation- π interactions, charge transfer interactions, entropic or hydrophobic or solvophobic effects.

15 A calix[n]pyrrole macrocycle attached to a solid support where n is 4, 5, 6, 7, or 8 is a particularly useful embodiment of the present invention. Due to the effective and selective separation of a variety of molecules by calix[n]pyrrole-derivatized solid supports as described in the examples herein, the present inventors have developed a new type of liquid phase chromatography termed "Hydrogen-Bonding Liquid Chromatography." The basis of the chromatography are noncovalent interactions, primarily hydrogen bonding.

20 A calix[n]pyrrole macrocycle having a first conformation in a solid state when unbound to a molecular or anionic species and having a second conformation in a solid state when bound to a molecular or anionic species where n is 4, 5, 6, 7, or 8 is another embodiment of the present invention. In particular, the present inventors have observed an unbound 1,3-alternate "first" conformation, and a bound cone or 1,2-alternate "second" conformation for calix[4]pyrrole macrocycles.

25 Exemplary calix[n]pyrrole macrocycles for anion and neutral molecular species binding, for solid supports, for derivatives, for multimers, and for conjugates, have structure I.





For structure I, when n is 4, $p = q = r = s = 0$, $R_1 - R_{16}$ are independently substituents as listed in paragraph i) below, and $R_A - R_D$ are independently substituents as listed in paragraph ii) below. When n is 5, $p = 1$, $q = r = s = 0$, R_1 to R_{20} are independently substituents as listed in paragraph i) below, and $R_A - R_E$ are independently substituents as listed in paragraph ii) below. When n is 6, $p = q = 1$, $r = s = 0$, R_1 to R_{24} are independently substituents as listed in paragraph i) below, and $R_A - R_F$ are independently substituents as listed in paragraph ii) below. When n is 7, $p = q = r = 1$, $s = 0$, R_1 to R_{28} are independently substituents as listed in paragraph i) below, and $R_A - R_G$ are independently substituents as listed in paragraph ii) below. When n is 8, $p = q = r = s = 1$, R_1 to R_{32} are independently substituents as listed in paragraph i) below, and $R_A - R_H$ are independently substituents as listed in paragraph ii) below;

i) hydrogen, halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxamide, carboxamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran,


tetrahydrothiapyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;

5 ii) hydrogen, alkyl, aminoalkyl, carboxy alkyl, carboxyamide alkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl.


or at least two substituents attached to any of the pyrrole β -carbons, *meso*-carbons or pyrrole nitrogens are coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein
10 in paragraph i) or ii) other than for bridged substituents;

Odd-numbered R-substituents are other than hydrogen for structure I.

New calix[n]pyrrole macrocycles are provided by the present invention having structure I as described above where alkyl is other than methyl or ethyl, where haloalkyl is other than chloropropyl, and wherein the calix[n]pyrrole is other than *meso*-tetramethyl-*meso*-
15 tetraphenylcalix[4]pyrrole, *meso*-octacyanopropylcalix[4]pyrrole, octaethyl-xanthoporphinogen, 5, 10, 15, 20-*meso*-tetrakis(3,5-Di-*tert*-butyl-4-quinomethide)porphyrinogen, or *meso*-octa-3-chloropropylcalix[4]pyrrole.



A calix[m]pyridino[n]pyrrole macrocycle having structure II is a further aspect of the present invention.




The term “m” designates a number of pyridines, and the term “n” designates a number of pyrroles in macrocycle II, $m+n=4$, and m is other than 1 or 2. When m is 4; $n=0$, $p=1$, and $q=0$, R101 to R108 and R201 to R212 are independently substituents as listed in paragraph i) above, and R301 - R304 are independently substituents as listed in paragraph ii) above. When m is 3; $n=1$, $p=0$, $q=1$, R101 to R108 and R201 to R211 are independently substituents as listed in paragraph i) above, and R301 - R304 are independently substituents as listed in paragraph ii) above. R201-R208 are other than hydrogen. When R301-R304 are other than a lone pair of electrons, $z=1$, that is the pyridine nitrogen atom is positively charged. When R301-R304 are a lone pair of electrons $z=0$, that is the pyridine nitrogen atom is neutral.

Further aspects of the present invention include calix[m]pyridino[n]pyrrole macrocycles where m and n designate a number of pyridines and pyrroles in the macrocycle, respectively, m and n are other than 0, and $m+n=5, 6, 7$, or 8. Each pyridine or pyrrole α -carbon is bound to another pyridine or pyrrole α -carbon via one non hydrogen-linked sp^3 -hybridized meso carbon; each sp^3 -hybridized meso carbon is further independently bonded to a group characterized in paragraph i) above with the exception of hydrogen; each pyridine β carbon, pyrrole β carbon and pyridine γ carbon is independently bonded to a group characterized in paragraph i) above; each pyridine or pyrrole nitrogen is bound to a group characterized in paragraph ii) above. In a further embodiment, at least one sp^3 meso carbon, pyridine β -carbon,

pyrrole β -carbon, pyridine γ carbon, or pyrrole nitrogen is coupled to form a bridged structure to itself or to another sp^3 hybridized *meso* carbon, pyridine β -carbon, pyrrole β -carbon, pyridine γ carbon, or pyrrole nitrogen; and when coupled to form a bridged structure, non-bridged atoms are as defined for an sp^3 hybridized *meso* carbon, pyridine β -carbon, pyrrole β -carbon, pyridine γ carbon, pyrrole nitrogen, or pyridine nitrogen.

A calix[m]pyridine macrocycle having structure III wherein m is 4, 5, 6, 7, or 8 is a further aspect of the present invention.



When m is 4, $p = q = r = s = 0$, R101 to R112 and R201 to R208 are independently substituents as listed in paragraph i) below, and R301 - R304 are independently substituents as listed in paragraph ii) above. When m is 5, $p = 1, q = r = s = 0$, R101 to R115 and R201 to R210 are independently substituents as listed in paragraph i) below, and R301 - R305 are independently substituents as listed in paragraph ii) above. When m is 6, $p = q = 1, r = s = 0$, R101 to R118 and R201 to R212 are independently substituents as listed in paragraph

i) below, and R₃₀₁ - R₃₀₆ are independently substituents as listed in paragraph ii) above. When m is 7, p=q=r= 1, s=0, R₁₀₁ to R₁₂₁ and R₂₀₁ to R₂₁₄ are independently substituents as listed in paragraph i) below, and R₃₀₁ - R₃₀₇ are independently substituents as listed in paragraph ii) above. When m is 8, p = q = r = s = 1, R₁₀₁ to R₁₂₄ and R₂₀₁ to R₂₁₆ are independently substituents as listed in paragraph i) below, and R₃₀₁ - R₃₀₈ are independently substituents as listed in paragraph ii) above. R₂₀₁-R₂₁₆ are other than hydrogen. When R₃₀₁-R_{30m} are other than a lone pair of electrons, z = 1, that is the pyridine nitrogen atom carries a positive charge. When R₃₀₁-R_{30m} are a lone pair of electrons z = 0, that is the pyridine nitrogen atom is neutral.

10

A calix[n]pyrrole where n is 5, 6, 7, or 8 is another embodiment of the present invention. Calix[n]pyrroles having a number of pyrrole rings greater than 4 have not been previously known.

The invention provides a number of different novel methods of making calix[n]pyrrole, calix[m]pyridino[n]pyrrole, and calix[m]pyridine macrocycles. A first method of making embodiment or synthesis embodiment is a method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having different substituents at *meso*-carbon atoms, comprising reacting pyrrole and at least two ketone molecules to produce a calix[n]pyrrole having different substituents at *meso*-carbon atoms.

A second method of making embodiment is a method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having an ester substituent, comprising reacting an ester-functionalized ketone, pyrrole and a ketone to produce a calix[n]pyrrole having an ester substituent.

A third method of making embodiment is a method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having an acid substituent, comprising reacting an ester-functionalized ketone, pyrrole and a ketone to produce a calix[n]pyrrole ester derivative; and de-esterifying the ester derivative to produce a calix[n]pyrrole monoacid derivative.

A method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising activating a calix[n]pyrrole where n is 4, 5, 6, 7, or 8; and reacting the activated calix[n]pyrrole with an electrophile is a fourth method of making embodiment of the invention.

A fifth synthetic method embodiment of the invention is a method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising reacting a β -substituted pyrrole and a ketone to produce a β -substituted calix[n]pyrrole.

A method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising producing a calix[n]pyrrole radical where n is 4, 5, 6, 7, or 8; and reacting the radical with a radical acceptor molecule is a sixth synthetic method embodiment of the invention.

A method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, 8 comprising reacting a pyrrole and a ketone in the presence of a Lewis acid is an seventh synthetic method embodiment of the invention.

5 A eighth method of making embodiment is a method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising reacting a pyrrole and a ketone in the presence of a heterogenous catalyst.

A ninth synthetic method embodiment of the invention is a method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising the steps of a) reacting a ketone-functionalized pyrrole and an alkyl metal to form a pyrrole alcohol, b) condensing the pyrrole alcohol to form a cyclizable pyrrole oligomer, and c) repeating step b) at least n minus 2 times
10 to form a calix[n]pyrrole.

An tenth synthetic method embodiment of the invention is a method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising reacting a pyrrole and a ketone in the presence of an anion temple.

15 A method of making a solid-supported calix[n]pyrrole, solid-supported calix[m]pyridino[n]pyrrole, or solid-supported calix[m]pyridine comprising attaching a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine having a functionalized group to a solid support reactive with the functionalized group is an eleventh synthetic method embodiment of the present invention.

20 A further composition of matter of the present invention is a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine made by one of the first through eleventh synthetic method embodiments provided herein.

A method of forming a complex of a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole, and an anion or a neutral molecule comprising contacting the calix[n]pyrrole or the
25 calix[m]pyridino[n]pyrrole with the anion or neutral molecule under conditions effective to allow the formation of the complex is a further aspect of the invention.

Another aspect of the invention is a method for separating a first molecule, anion or cation from a mixture of a first molecule, anion or cation and other species, comprising obtaining a calix[n]pyrrole-, a calix[m]pyridino[n]pyrrole- or a calix[m]pyridine-derivatized solid
30 support; and contacting the solid support with the mixture to separate the first molecule, anion or cation any other materials present. Another method of separating includes a batch processing where a macrocycle of the present invention is added to a mixture containing the molecule to be separated, and separating the macrocycle-molecule complex. Macrocycles provided herein are particularly effective as separation media as described in the present
35 examples.

A method of transporting a molecular or anionic species through a membrane comprising incorporating a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole into the membrane; and contacting the membrane with the molecular or anionic species in the presence of a gradient of

the molecular or anionic species or counter gradient of another back-transported species is an embodiment of the invention. Thus the present invention maybe used to facilitate the transport of anionic or neutral substrates in both a synport or antiport sense.

A further embodiment of the present invention is an electropolymerizable
 5 calix[n]pyrrole, calix[m]pyridino[n]pyrrole, or calix[m]pyridine.

A further embodiment of the present invention is an anion, cation, or neutral molecule ion-selective electrode comprising a conductive body, a polymer, and a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine.

A further embodiment of the present invention is an ion-selective electrode wherein the
 10 calix[n]pyrrole, the calix[m]pyridino[n]pyrrole, or the calix[m]pyridine is electropolymerized and forms the conductive body.

A method of electrochemical detection of an anion, a cation, or a neutral molecule comprising assembling the ion selective electrode and contacting the electrode with a solution of the anion, the cation, or the neutral molecule and detecting the presence or absence of the anion,
 15 the cation, or the neutral molecule is an embodiment of the present invention.

A method of binding a cation comprising contacting the cation with a calix[n]pyrrole or calix[m]pyridino[n]pyrrole having a cation-binding functionality, or with a calix[m]pyridine is also an aspect of the invention.

Further embodiments include a chromatography column or a sensor comprising a solid
 20 support bound to a calix[n]pyrrole where n is 4-8, to a calix[m]pyridino[n]pyrrole where $m+n=4-8$ and m is not 1 or 2, or to a calix[m]pyridine where m is 4-8. Use of a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine in the preparation of a pharmaceutical composition for use in *in vivo* or *ex vivo* treatment of body tissues is another embodiment of the invention. The treatment may involve binding, transport, and/or removal of ions or neutral
 25 molecular species for conditions such as gout, for kidney dialysis, for removal of viruses, for introduction of antiviral drugs, or the like. A method therefore includes administering to a patient in need thereof a therapeutically effective amount of a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine.

30 BRIEF DESCRIPTION OF THE DRAWINGS.

FIG. 1A illustrates the X-ray crystal structure of octamethylcalix[4]pyrrole, 1, adopting a 1,3-alternate conformation. Thermal ellipsoids are at 30% probability level.

FIG. 1B illustrates the X-ray crystal structure of *meso*-tetraspirocyclohexylcalix[4]pyrrole, 2, adopting a 1,3-alternate conformation. Thermal
 35 ellipsoids are at 30% probability level.

FIG. 2A illustrates the X-ray crystal structure of *meso*-octamethylcalix[4]pyrrole tetra-butylammonium chloride complex, 1-Cl⁻. The *meso*-octamethylcalix[4]pyrrole adopts a cone

conformation wherein all four pyrrole NH groups are hydrogen-bonded to the bound chloride anion. The counter cation is omitted for clarity. Thermal ellipsoids are at 30% probability level.

FIG. 2B illustrates the X-ray crystal structure of *meso*-tetraspirocyclohexylcalix[4]pyrrole tetrabutylammonium chloride complex, 2·Cl⁻. The *meso*-tetraspirocyclohexylcalix[4]pyrrole adopts a cone conformation wherein all four pyrrole NH groups are hydrogen-bonded to the bound chloride anion. The counter cation is omitted for clarity. Thermal ellipsoids are at 30% probability level.

FIG. 3A illustrates the X-ray crystal structure of *meso*-octamethylcalix[4]pyrrole bis-methanol complex, 1·2 MeOH. The *meso*-octamethylcalix[4]pyrrole adopts a 1,3-alternate conformation wherein two NH groups bind to each methanol. Thermal ellipsoids are at 30% probability level.

FIG. 3B illustrates the X-ray crystal structure of *meso*-octamethylcalix[4]pyrrole bis-dimethylformamide complex, 1·2 DMF. The *meso*-octamethylcalix[4]pyrrole adopts a 1,2-alternate conformation wherein two NH groups bind to each dimethylformamide. Thermal ellipsoids are at 30% probability level.

FIG. 3C illustrates the X-ray crystal structure of *meso*-octamethylcalix[4]pyrrole dimethylsulfoxide complex, 1·DMSO. The *meso*-octamethylcalix[4]pyrrole adopts a 1,3-alternate conformation wherein one NH group binds to a dimethylsulfoxide. Thermal ellipsoids are at 30% probability level.

FIG. 4 illustrates the X-ray crystal structure of tetrabutylammonium salt of calix[4]pyrrole, 4. Calix[4]pyrrole anions self-assemble in the solid state forming dimers. Thermal ellipsoids are at 30% probability level. Counter cations are omitted for clarity.

FIG. 5 illustrates the X-ray crystal structure of a *meso*-octabenzylcalix[4]pyrrole acetone complex, 6·3(acetone). The octabenzylcalix[4]pyrrole adopts a 1,3-alternate conformation wherein one NH group binds to an acetone. An unbound acetone molecule in the crystal lattice is omitted for clarity. Thermal ellipsoids are at 30% probability level.

FIG. 6 illustrates the X-ray crystal structure of *meso*-tetrakis-tetrahydrothiopyranylcalix[4]pyrrole, 7, adopting a 1,3-alternate conformation. Thermal ellipsoids are at 30% probability level.

FIG. 7 illustrates an X-ray crystal structure of β-octabromo-*meso*-octamethylcalix[4]pyrrole, 14, adopting a 1,3-alternate conformation. Thermal ellipsoids are at 30% probability level.

FIG. 8 illustrates an X-ray crystal structure of a *meso*-tetramethylcalix[4]pyrrole-*p*-*tert*-butylcalix[4]arene pseudo dimer, 15. Thermal ellipsoids are at 30% probability level.

FIG. 9 illustrates a ¹H NMR of a *meso*-tetramethylcalix[4]pyrrole-*p*-*tert*-butylcalix[4]arene pseudo dimer in CD₂Cl₂.

FIG. 10 illustrates a sodium chloride complex of calix[4]pyrrole-crown ether conjugate, 32.

FIG. 11A illustrates a cyclic voltammogram of compound, **33**, conducted in dichloromethane at a scan rate of 100 mV/sec. Working electrode: platinum disk, counter electrode: platinum wire, reference electrode: Ag/AgCl. 0.1 M tetrabutylammonium hexafluorophosphate as base electrolyte. Scale equals 2 microamps.

5 FIG. 11B illustrates a cyclic voltammogram of compound, **34**, conducted in dichloromethane at a scan rate of 100 mV/sec. Working electrode: platinum disk, counter electrode: platinum wire, reference electrode: Ag/AgCl. 0.1 M tetrabutylammonium hexafluorophosphate as base electrolyte. Scale equals 2 microamps.

10 FIG 12A illustrates a cyclic voltammogram of solutions of compound, **33**, and 2 equivalents of tetrabutylammonium fluoride (1), tetrabutylammonium chloride (2), and tetrabutylammonium dihydrogenphosphate (3) conducted in dichloromethane at a scan rate of 100 mV/sec. Working electrode: platinum disk, counter electrode: platinum wire, reference electrode: Ag/AgCl. 0.1 M tetrabutylammonium hexafluorophosphate as base electrolyte. Scale equals 5 microamps.

15 FIG 12B illustrates a cyclic voltammogram of solutions of compound, **34**, and 2 equivalents of tetrabutylammonium fluoride (1), tetrabutylammonium chloride (2), and tetrabutylammonium dihydrogenphosphate (3) conducted in dichloromethane at a scan rate of 1200 mV/sec. Working electrode: platinum disk, counter electrode: platinum wire, reference electrode: Ag/AgCl. 0.1 M tetrabutylammonium hexafluorophosphate as base electrolyte. Scale equals 5 microamps.

20 FIG. 13 illustrates a calix[4]pyrrole-R-(+)- α -methylbenzylamine conjugate, **40**, -fluoride complex represented as a diastereomeric mixture.

FIG. 14 illustrates the hydrogen bond donating capabilities of a mono- β -butylamidocalix[4]pyrrole, **28**, complexed to an anion.

25 FIG. 15A illustrates *meso*-substituted-calix[4]pyrrole substituted silica gel, **M**.

FIG. 15B illustrates β -substituted-calix[4]pyrrole substituted silica gel, **B**.

FIG. 16 illustrates the separation of phenyl arsenate (1), benzene sulfonate (2), benzoate (3), and phenyl phosphate (4) on a *meso*-substituted calix[4]pyrrole column: mobile phase, 50 mM phosphate buffer, pH = 7.0; flow rate 0.3 mL/min.; column temperature 25°C; detection 254 nm (0.100 AUFS).

30 FIG. 17 illustrates the separation of 5'-adenosine monophosphate (1), 5'-adenosine diphosphate (2), and 5'-adenosine triphosphate (3) on a *meso*-substituted calix[4]pyrrole column: mobile phase 250 mM phosphate buffer, pH = 7.0; flow rate 0.3 mL/min.; column temperature 25°C; detection 254 nm (0.100 AUFS).

35 FIG. 18 illustrates the separation of 5'-adenosine monophosphate (1), 5'-adenosine diphosphate (2), and 5'-adenosine triphosphate (3) on a β -substituted calix[4]pyrrole column: mobile phase 105 mM phosphate buffer, pH = 7.0; flow rate 0.20 mL/min.; column temperature 25°C; detection UV at 254 nm (0.100 AUFS).

FIG. 19 illustrates the separation of uridine (1), phenol (2), aniline (3), benzene (4), and 4 iodo-nitro benzene (5) on a *meso*-substituted calix[4]pyrrole column: mobile phase 60% water/ 40% acetonitrile (v/v); flow rate 0.3 mL/min.; column temperature 25°C; detection 254 nm (0.100 AUFS).

5 FIG. 20 illustrates the separation of Cbz-N-protected- serine (1), glutamine (2), alanine (3), phenylalanine (4), tryptophan (5), aspartate (6), and glutamate (7) on a *meso*-substituted calix[4]pyrrole column: mobile phase 70% 30 mM acetate buffer, pH = 7.0/ 30 % CH₃CN (v/v); flow rate 0.3 mL/min.; column temperature 25°C, detection 254 nm (0.100 AUFS).

10 FIG. 21 illustrates the separation of Cbz-N-protected- serine (1), leucine (2), alanine (3), phenylalanine (4), tryptophan (5), aspartate (6), and glutamate (7) on a β -substituted calix[4]pyrrole column: mobile phase 75/25 30 mM acetate buffer, pH = 7.0/ acetonitrile; flow rate 0.20 mL/min.; column temperature 25°C; detection UV at 254 nm (0.100 AUFS).

15 FIG. 22 illustrates the separation of 4-fluorobiphenyl (1), 4,4'-difluorobiphenyl (2), 2,2',3,3',5,5',6,6',-octafluorobiphenyl (3), and perfluorobiphenyl (4) on a *meso*-substituted calix[4]pyrrole column: mobile 60% water/40% acetonitrile (v/v); flow rate 0.3 mL/min.; column temperature 25°C; detection 254 nm (0.100 AUFS).

20 FIG. 23 illustrates the separation of 4-fluorobiphenyl (1), 4,4'-difluorobiphenyl (2), 2,2',3,3',5,5',6,6',-octafluorobiphenyl (3), and perfluorobiphenyl (4), on a β -substituted calix[4]pyrrole column: mobile 76% water/24% acetonitrile (v/v); flow rate 0.2 mL/min.; column temperature 25°C; detection 254 nm (0.100 AUFS).

25 FIG. 24 illustrates the separation of a mixture of polydeoxythymidilates oligomers containing between (12) and (18) base pairs on a *meso*-substituted calix[4]pyrrole column: mobile phase 50/50 acetonitrile/ 50 mM phosphate, 250 mM chloride buffer, pH = 6.8 (v/v); flow rate 0.4 mL/min.; column temperature 25°C; detection UV at 265 nm.

30 FIG. 25 illustrates the separation of a mixture of polydeoxythymidilates oligomers containing between (12) and (18) base pairs on a β -substituted calix[4]pyrrole column: mobile phase 40/60 acetonitrile/ 40 mM phosphate, 50 mM chloride buffer, pH = 7.0 (v/v), flow rate 0.25 mL/min.; column temperature 25°C; detection UV at 262 nm (0.100 AUFS).

35 FIG. 26 illustrates the separation of TCTAGA, (1), GCATGC, (2), and CCCGGG, (3), on a *meso*-substituted calix[4]pyrrole column: mobile phase 50/50 acetonitrile/ 50 mM sodium phosphate, pH = 7.0 (v/v); flow rate 0.4 mL/min.; column temperature 25°C; detection UV at 265 nm (0.100 AUFS).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS


35 The present invention provides calix[n]pyrroles, calix[m]pyridines, calix[m]pyridino[n]pyrroles, calix[n]pyrrole-anion complexes, calix[m]pyridino[n]pyrrole-anion complexes, calix[n]pyrrole-neutral molecule complexes, calix[m]pyridino[n]pyrrole-neutral molecule complex, calix[m]pyridine-cation complexes, calix[m]pyridine-neutral molecule

complexes and calix[n]pyrroles having a first conformation in a solid-state when unbound and having a second conformation when non-covalently bound to a guest species, as new compositions of matter.


Specific anions include, but are not limited to, halide anions, carboxylates, phosphates, sulfates, oxalates, terephthalates, phospholipids, nucleotides, oligonucleotides, DNA, RNA, or anionic polyoxometallates such as pertechnetate.

The term "molecular species" as used herein, means a neutral molecule, and represents a variety of classes of molecules since the macrocycles of the present invention provide different sizes of cavities and varieties of donor and acceptor sites. Specific neutral molecules include, but are not limited to, alcohols, polyalcohols, ketones, polyketones, phenols, polyhydroxylated aliphatic and aromatic compounds, amino compounds, amino acids, urea, guanidine, saccharides, biologically important polymers like proteins derivatives and the like.

Carbon atoms in pyrrole rings are referred to as α or β , α carbons are adjacent to the NH group and β -carbons are adjacent to the α -carbons (see IV). An alternative designation is used for all the atoms in the pyrrole ring. The nitrogen atom is labelled 1 and the other atoms are numbered sequentially starting from an adjacent carbon atom. Thus the α -carbons are also referred to as atoms 2 and 5 and the β -carbons as atoms 3 and 4.



IV



V

Similarly, carbon atoms in the pyridine rings are referred to as α , β or γ , α carbons are adjacent to the N atom, β -carbons are adjacent to the α -carbons and the γ -carbon is adjacent to the two β -carbons (see V). An alternative designation is used for all the atoms in the pyridine ring. The nitrogen atom is labelled 1 and the other atoms are numbered sequentially starting from an adjacent carbon atom. Thus α -carbons are also referred to as atoms 2 and 6, β -carbons as atoms 3 and 5, and the γ -carbon as atom 4.

Substituents for Macrocycles of the Present Invention

Representative examples of alkanes useful as alkyl group substituents of the present invention include straight-chain, branched or cyclic isomers of propane, butane, pentane, hexane, heptane, octane, nonane and decane, with methane, ethane and propane being preferred. Alkyl groups having up to about thirty, or up to about fifty carbon atoms are contemplated in

the present invention. Representative examples of substituted alkyls include alkyls substituted by two or more functional groups as described herein.

Representative examples of alkenes useful as alkenyl group substituents include ethene, straight-chain, branched or cyclic isomers of propene, butene, pentene, hexene, heptene, octene, nonene and decene, with ethene and propene being preferred. Alkenyl groups having up to about thirty or fifty carbon atoms, and up to about five double bonds, or more preferably, up to about three double bonds are contemplated in the present invention.

Representative examples of alkynes useful as alkynyl group substituents include ethyne, straight-chain, branched or cyclic isomers of propyne, butyne, pentyne, hexyne, heptyne, octyne, nonyne and decyne, with ethyne and propyne being preferred. Alkynyl groups having up to about thirty, or up to about fifty carbon atoms, and having up to about five or up to about three triple bonds are contemplated in the present invention.

The aryl may be a compound whose molecules have the ring structure characteristic of benzene, naphthalene, phenanthrene, anthracene, and the like, i.e., either the 6-carbon ring of benzene or the condensed 6-carbon rings of the other aromatic derivatives. For example, an aryl group may be phenyl or naphthyl, and the terms used herein include both unsubstituted aryls and aryls substituted with one or more nitro, carboxy, sulfonic acid, hydroxy, oxyalkyl or halide substituent. Purine or pyrimidine molecules are included as "aryl" molecules. In this case, the substituent on the phenyl or naphthyl may be added in a synthetic step after the condensation step which forms the macrocycle.

Among the halide substituents, chloride, bromide, fluoride and iodide are contemplated in the practice of this invention. Representative examples of haloalkyls used in this invention include halides of methane, ethane, propane, butane, pentane, hexane, heptane, octane, nonane and decane, with halides, preferably chlorides or bromides, of methane, ethane and propane being preferred.

"Hydroxyalkyl" means alcohols of alkyl groups. Preferred are hydroxyalkyl groups having one to twenty, more preferably one to ten, hydroxyls. "Hydroxyalkyl" is meant to include glycols and polyglycols; diols of alkyls, with diols of C1-10 alkyls being preferred, and diols of C1-3 alkyls being more preferred; and polyethylene glycol, polypropylene glycol and polybutylene glycol as well as polyalkylene glycols containing combinations of ethylene, propylene and butylene.

Representative examples of oxyalkyls include the alkyl groups as herein described having ether linkages. "Oxyalkyl" is meant to include polyethers with one or more functional groups. The number of repeating oxyalkyls within a substituent may be up to 200, preferably is from 1-20, and more preferably, is 1-10, and most preferably is 1-5. A preferred oxyalkyl is $O(CH_2CH_2O)_xCH_3$ where $x = 1-100$, preferably 1-10, and more preferably, 1-5.

"Oxyhydroxyalkyl" means alkyl groups having ether or ester linkages, hydroxyl groups, substituted hydroxyl groups, carboxyl groups, substituted carboxyl groups or the like.

Representative examples of thioalkyls include thiols of ethane, thiols of straight-chain, branched or cyclic isomers of propane, butane, pentane, hexane, heptane, octane, nonane and decane, with thiols of ethane (ethanethiol, C_2H_5SH) or propane (propanethiol, C_3H_7SH) being preferred. Sulfate-substituted alkyls include alkyls as described above substituted by one or more sulfate groups, a representative example of which is diethyl sulfate ($(C_2H_5)_2SO_4$).

Representative examples of phosphates include phosphate or polyphosphate groups. Representative examples of phosphate-substituted alkyls include alkyls as described above substituted by one or more phosphate or polyphosphate groups. Representative examples of phosphonate-substituted alkyls include alkyls as described above substituted by one or more phosphonate groups.

Representative examples of carboxy groups include carboxylic acids of the alkyls described above as well as aryl carboxylic acids such as benzoic acid and derivatives thereof. Representative examples of carboxyamides include primary carboxyamides ($CONH_2$), secondary ($CONHR'$) and tertiary ($CONR'R''$) carboxyamides where each of R' and R'' is a functional group as described herein.

Representative examples of useful amines include a primary, secondary or tertiary amine of an alkyl as described herein.

"Carboxyamidealkyl" means alkyl groups with secondary or tertiary amide linkages or the like. "Carboxyalkyl" means alkyl groups having hydroxyl groups, carboxyl or amide substituted ethers, ester linkages, tertiary amide linkages removed from the ether or the like.

The term "saccharide" includes oxidized, reduced or substituted saccharide; hexoses such as D-glucose, D-mannose or D-galactose; pentoses such as D-ribose or D-arabinose; ketoses such as D-ribulose or D-fructose; disaccharides such as sucrose, lactose, or maltose; derivatives such as acetals, amines, and phosphorylated sugars; oligosaccharides, as well as open chain forms of various sugars, and the like. Examples of amine-derivatized sugars are galactosamine, glucosamine, sialic acid and D-glucamine derivatives such as 1-amino-1-deoxysorbitol.

By 'reporter group' is meant a substituent that is fluorescent, chromophoric, electropolymerizable, redox-active, or optically active. Examples of a fluorescent reporter group include, but are not limited to, ruthenium (II) bipyridyl complexes, acetylnaphthalene, 9-aminoacridine, 9-phenylanthracene, benzimidazole, *N*-methylbenzo[b]carbazole, 2-phenylbenzoxazole, 1,1'-binaphthyl, fluorene, fluorescein dianion, indeno[2, 1-a]indene, 2,5-diphenylfuran, perylene, 2-aminopurine, *p*-quaterphenyl, 4,4'-diphenylstilbene, sapphyrins and texaphyrins. Examples of a chromophoric reporter groups include isosulfan blue, fluoroscein, 2',7'-dichlorofluorescein, rhodamine, carboxyrhodamine 6G, dialkylaminocoumarin, erythrosin, pyrene, Texas Red, Rhodol Green, Malachite Green, 4,4'-difluoro-5,7-dimethyl-4-bora-3a,4a,diaza-5-indacene-3-propionic acid, Oregon Green 488, 9-(diethylamino)-5-octadecanoylimino-5H-benzo[a]phenoxazine (Chromoionophore I), 9-dimethylamino-5-[4-(16-butyl-2,14-dioxo-3,15-dioxaeicosyl)phenylimino]benzo[a]phenoxazine, 9-dimethylamino-5-[4-

- (16 (Chromoionophore II), 9-(diethylamino)-5,9-(di[(2-octyldecyl)imino]benzo[a]phenoxazine (Chromoionophore III), 5-octadecanoyloxy-2-(4-nitrophenylazo)phenol (Chromoionophore IV), 9-(diethylamino)-5-(2-naphthoylimino)-5H-benzo[a]phenoxazine (Chromoionophore V), 4',5'-dibromofluorescein octadecyl ester (Chromoionophore VI), Chromomycin A₃, 2-(4-nitrophenylazo)chromotropic acid disodium salt (Chromotrop 2B), 2-(phenylazo)chromotropic acid disodium salt (Chromotrop 2 R), 4,5-dihydroxynaphthalene-2,7-disulfonic acid disodium salt (Chromotropic acid disodium salt dihydrate), Cibacron Blue F3G-A (Procion Blue H-B; Reactive Blue 2), Chromeazurol B (C.I.No. 43830), Chromeazurol S (C.I.No. 43825), 5,7-dihydroxyflavone, 5,7-dinitro-8-hydroxy-2-naphthalenesulphonic acid (Naphthol Yellow S), 4,4-dimethylaminoazobenzene (Dimethyl Yellow), *N*-ethyl-*N*-[4-[[4-[ethyl[(3-sulfophenyl)methyl]amino]phenyl](4-sulfophenyl)-methylene]2,5-cyclohexadien-1-ylidene]-3-sulfobenzenemethanaminium hydroxide, inner salt, disodium salt (Light Green SF Yellowish, Acid Green), *N*-ethyl-*N*-[4-[[4-[ethyl[(3-sulfophenyl)methyl]amino]phenyl](4-hydroxy-2-sulfophenyl)-methylene]2,5-cyclohexadien-1-ylidene]-3-sulfobenzenemethanaminium hydroxide, inner salt, disodium salt (Fast Green FCF), 6,6'-[(3,3'-dimethyl[1,1'-biphenyl]4,4'-diyl)bis(azo)]bis[4-amino-5-hydroxy-1,3-naphthalenedisulfonic acid] tetrasodium salt (Evan's Blue), 4,5,6,7,-tetrachloro-3',6'-dihydroxy-2',4',5',7'-tetraiodospiro[isobenzofuran-1(3H),9'-[9H]xanthen]-3-one dipotassium salt (Rose Bengal), 3,7-bis(dimethylamino)phenothiazin-5-ium chloride (Methylene Blue), 3,6-bis(dimethylamino)acridine hydrochloride zinc chloride double salt (Acridine Orange), 2',4',5',7'-tetrabromo-3',6'-dihydroxyspiro[isobenzofuran-1(3H),9'-[9H]xanthen]-3-one disodium salt (Eosine Yellowish), 4',5'-dibromo-3',6'-dihydroxy-2',7'-dinitrospiro[isobenzofuran-1(3H),9'-[9H]xanthen]-3-one disodium salt (Eosine I Bluish), sapphyrins and texaphyrins. By "electropolymerizable" is meant a moiety that will polymerize when subjected to a particular voltammetric potential or a continuous scanning potential. Examples of an electropolymerizable group include, but are not limited to, α -free pyrroles, α -free thiophenes or vinyl groups. By "redox-active" is meant a moiety which can undergo an oxidation or a reduction process. Examples of a redox-active group include, but are not limited to, ferrocene, cobaltocenium, ruthenium (II) bipyridyl complexes, transition metals, fullerenes, porphyrins, expanded porphyrins and pyrroles.
- In one embodiment of the present invention, a calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine is further coupled to a site-directing molecule. "Site-directing" means having specificity for targeted sites. "Specificity for targeted sites" means that upon contacting the calix[n]pyrrole-, calix[m]pyridino[n]pyrrole- or calix[m]pyridine-conjugate with the targeted site, for example, under physiological conditions of ionic strength, temperature, pH and the like, specific binding will occur. The interaction may occur due to specific electrostatic, hydrophobic, or other interaction of certain residues of the conjugate with specific residues of the target to form a stable complex under conditions effective to promote the interaction.

Exemplary site-directing molecules contemplated in the present invention include but are not limited to: polydeoxyribonucleotides; oligodeoxyribonucleotides; polyamides, including peptides having affinity for a biological receptor, and proteins such as antibodies; steroids and steroid derivatives; hormones such as estradiol, or histamine; hormone mimics such as morphine; and further macrocycles such as sapphyrins and rubyrins. A preferred site-directing molecule is an oligonucleotide.

Representative examples of useful steroids include a steroid hormone of the following five categories: progestins (e.g. progesterone), glucocorticoids (e.g., cortisol), mineralocorticoids (e.g., aldosterone), androgens (e.g., testosterone) and estrogens (e.g., estradiol).

The term "a peptide having affinity for a biological receptor" means that upon contacting the peptide with the biological receptor, for example, under appropriate conditions of ionic strength, temperature, pH and the like, specific binding will occur. The interaction may occur due to specific electrostatic, hydrophobic, entropic or other interaction of certain amino acid or glycolytic residues of the peptide with specific amino acid or glycolytic residues of the receptor to form a stable complex under the conditions effective to promote the interaction. The interaction may alter the three-dimensional conformation and the function or activity of either or both the peptide and the receptor involved in the interaction. A peptide having affinity for a biological receptor may include an endorphin, an enkephalin, a growth factor, e.g. epidermal growth factor, poly-L-lysine, a hormone, a peptide region of a protein and the like. A hormone may be estradiol, for example.

Representative examples of useful amino acids of peptides or polypeptides include amino acids with simple aliphatic side chains (e.g., glycine, alanine, valine, leucine, and isoleucine), amino acids with aromatic side chains (e.g., phenylalanine, tryptophan, tyrosine, and histidine), amino acids with oxygen and sulfur-containing side chains (e.g., serine, threonine, methionine, and cysteine), amino acids with side chains containing carboxylic acid or amide groups (e.g., aspartic acid, glutamic acid, asparagine, and glutamine), and amino acids with side chains containing strongly basic groups (e.g., lysine and arginine), and proline. Representative examples of useful peptides include any of both naturally occurring and synthetic di-, tri-, tetra-, pentapeptides or longer peptides derived from any of the above described amino acids (e.g., endorphin, enkephalin, epidermal growth factor, poly-L-lysine, or a hormone). Representative examples of useful polypeptides include both naturally occurring and synthetic polypeptides (e.g., insulin, ribonuclease, and endorphins) derived from the above described amino acids and peptides.

The term "binding agent" means that upon contacting the binding agent with a guest species for binding under appropriate conditions of ionic strength, temperature, pH and the like, specific binding will occur. The interaction may alter the three-dimensional conformation and the function or activity of either or both the bound species and the receptor involved in the interaction. Preferred binding agents include a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, a

calix[m]pyridine, a calixarene, a cation-binding functionality, a crown ether, a chelating group, a porphyrin, or an expanded porphyrin such as sapphyrin, rosarin, rubyrin, texaphyrin, amethyryn or turcasarin, or oligopyrroles.

A "catalytic group," as used herein, means a chemical functional group that may act as a general acid, Brønsted acid, general base, Brønsted base, nucleophile, or any other means by which an activation barrier to reaction is lowered or the ground state energy of a substrate is increased. Exemplary catalytic groups contemplated include, but are not limited to, imidazole; guanidine; zinc coordinated to a nitrogen containing macrocycle, EDTA complexes, DTPA complexes, substituted saccharides such as a D-glucosamine, D-mannosamine, D-galactosamine, D-glucamine, and the like; amino acids such as L-histidine and L-arginine; derivatives of amino acids such as histamine; polymers of amino acids such as poly-L-lysine, (LysAla)_n, (LysLeuAla)_n where n is from 1-30 or preferably 1-10 or more preferably 2-7 and the like. The catalytic group may be attached either directly to the calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine or *via* a linker or couple of variable length.

A couple may be described as a linker, i.e., the covalent product formed by reaction of a reactive group designed to attach covalently another molecule at a distance from the calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine macrocycle. Exemplary linkers or couples are amides, amine, disulfide, thioether, ether, ester, or phosphate covalent bonds.

In most preferred embodiments, conjugates and appended groups are covalently bonded to the calix[n]pyrrole *via* a carbon-carbon, carbon-nitrogen, carbon-sulfur, or a carbon-oxygen bond, more preferably a carbon-carbon, carbon-oxygen or a carbon-nitrogen bond.

In the presently preferred macrocycles, 3 - 20 and 22 - 42 are preferred. Most preferred are macrocycles 3, 4, 10, 12 and solid supported materials Gel M and Gel B. While the cited calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines are presently preferred for use in the present invention, the invention is not limited thereto and any calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine may be used.

Synthetic Methods for Macrocycles of the Present Invention

The present inventors have used a number of different macrocycles of the present invention. The following preferred embodiments are described.

Calix[n]pyrroles having different substituents at meso-carbons. A method of synthesizing calix[n]pyrroles having different substituents at *meso*-carbons comprises reacting a ketone with a pyrrole in the presence of a Brønsted acid. Examples of ketones include, but are not limited to, acetone, cyclohexanone, 3-pentanone, 5-heptanone, or dibenzylacetone. Examples of pyrroles include, but are not limited to, pyrrole, 3,4-dimethyl pyrrole, 3,4-diethyl pyrrole, 3,4-dimethoxypyrrole or 3-butylpyrrole. Examples of Brønsted acids include, but are not limited to HCl, methanesulfonic acid, toluenesulfonic acid, benzenesulfonic acid, HClO₄, formic acid, acetic acid, sulfuric acid or phosphoric acid.

template. Examples of pyrroles, ketones, Lewis acids, Brønsted acids, and heterogeneous catalysts have been given herein. Examples of templates include, but are not limited to, hydrogen bonding neutral molecules, such as sugars, cyclodextrins, alcohols or amines or anionic species such as, halides, carboxylates, sulfonates or phosphates.

5 Using halides to illustrate, but not limit, the methods of template synthesis of calix[n]pyrroles, pyrrole and acetone would be condensed using a Brønsted acid, a Lewis acid or a homogeneous catalyst in the presence of a halide. By varying the size of the halide, it would be possible to “tune” the size of calix[n]pyrrole where n is 4, 5, 6, 7, or 8.

10 One skilled in the art of organic synthesis in light of the present disclosure could extend and refine the referenced basic synthetic chemistry to produce calix[n]pyrroles having various substituents. For example, polyether-linked polyhydroxylated groups, saccharide substitutions in which the saccharide is appended via an acetal-like glycosidic linkage, an oligosaccharide or a polysaccharide may be similarly linked to a calix[n]pyrrole. A carboxylated calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine in which the acid groups are linked to the
15 calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine core *via* aryl ethers or functionalized alkyl substituents could be converted to various amido products wherein the amide linkages serve to append further substituents. Saccharide moieties may be appended *via* amide bonds.

Calixarene-calixpyrrole pseudo dimers, multimers and polymers

20 Calix[n]arenes are macrocyclic molecules composed of phenol rings linked by methylene groups. According to one aspect of the present invention, calix[4]pyrroles are useful as anion binding agents. A method of the invention comprises reacting a cyclic ketone-containing compound with pyrrole. The number of ketones and the number of pyrroles will be equal to 4. The cyclic ketone-containing compound acts as a template upon which the calixpyrrole is
25 formed, after which the synthesized calixpyrrole may be detached from the ketone-containing template. Alternatively, where the template is a calixarene, the resulting cylindrical calix[4]arene-calix[4]pyrrole pseudo dimer may be kept intact.

30 The cyclic ketone-containing compound may be selected from any cyclophane molecule having 4 ketone groups. Examples of such molecules include, but are not limited to, calix[4]arenes, resorcinarenes, CTV, and cyclic sugars such as the cyclodextrins. Alternatively, benzene rings substituted with ketone groups, or azacrown ethers with *N*-pendant ketone moieties may serve as templates in the synthesis of the present invention. In one aspect of the invention, it is preferred that the ketone groups of the template all be pointing in the same direction. For example, in the case of calixarenes, the calixarene is in a cone conformation.

35 In order to detach a templated calix[n]pyrrole, a cleavable group is present between the template and the ketone. This cleavable group may be covalently linked or non-covalently linked. The cleavage group may be, for example, an ester that can be subsequently hydrolyzed, or a benzyl group that can be hydrogenated or removed with a de-benzylating agent such as

Using acetone and cyclohexanone to illustrate this synthetic method embodiment, acetone and/or cyclohexanone react with pyrrole in the presence of a Brønsted acid to form a macrocyclic structure comprised of n pyrrole units, where n is 4, 5, 6, 7, or 8 and n substituted *meso*-carbons where n is 4, 5, 6, 7, or 8.

- 5 ***Calix[n]pyrroles having ester functionalities at meso-carbon(s).*** A method of making calix[n]pyrroles having ester functionalities at *meso*-carbon(s) comprises reacting a ketone having one or more ester functionalities with a pyrrole in the presence of a Brønsted acid. The method of invention additionally comprises reacting a mixture of ketones containing one or more ester functionalities and ketones not containing ester functionalities with a Brønsted acid to
10 provide a range of ester functionalities. Example of a ketone having one or more ester functionalities include, but are not limited to, methylacetylbutyrate, acetylpropionate and acetylpentanoate. Examples of pyrroles include, but are not limited to, pyrrole, 3,4-diethylpyrrole, 3,4-dimethoxypyrrole or 3-butylpyrrole. Examples of Brønsted acids include, but are not limited to HCl, methanesulfonic acid, toluenesulfonic acid, benzenesulfonic acid,
15 HClO₄, formic acid, acetic acid, sulfuric acid or phosphoric acid.

Using methylacetylbutyrate to illustrate the method of the present invention, methylacetylbutyrate reacts with pyrrole in the presence of a Brønsted acid to form a macrocyclic structure comprised of n pyrrole units, where n is 4, 5, 6, 7, or 8 and the number of ester functionalities is equal to n .

- 20 ***Calix[n]pyrroles having acid functionalities at the meso-carbon(s).*** This method embodiment for making calix[n]pyrroles having acid functionalities at the *meso*-carbons comprises de-esterifying an ester functionalized calix[n]pyrrole where n is 4, 5, 6, 7 or 8. De-esterifying is accomplished by heating the ester functionalized calix[n]pyrrole in a water/ethanol solution in the presence of sodium hydroxide and subsequent acidification and extraction of the
25 calixpyrrole acid.

- Addition of an electrophile to form β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8.*** This method embodiment comprises activating a calix[n]pyrrole, where n is 4, 5, 6, 7 or 8 and reacting the activated calix[n]pyrrole with an electrophile in order to make a β -substituted calix[n]pyrrole. Examples of activation include, but are not limited to activation with a
30 Brønsted base such as *n*-butyl lithium, *tert*-butyl lithium, phenyl lithium or any of these Brønsted bases described herein in conjunction with tetramethylethylenediamine. Examples of electrophiles include, but are not limited to, ethylbromoacetate, methylbromoacetate, acrylonitrile, methyl acrylate, acyl chloride, alkyl halide, functionalized alkyl halides or carbon dioxide.

- 35 An exemplary method is found in Example 3.

β -Substituted calix[n]pyrroles where n is 4, 5, 6, 7, or 8.

A method of making a β -substituted calix[n]pyrroles where n is 4, 5, 6, 7, or 8 comprises reacting a β -substituted pyrrole with a ketone in the presence of a Brønsted acid to produce a

calix[n]pyrrole where n is 4, 5, 6, 7 or 8. Examples of β -substituted pyrroles include, but are not limited to, 3,4-dimethoxypyrrole, 3,4-diethyl pyrrole 3,4-dimethylpyrrole, 3,4-diphenylpyrrole, or 3-butylpyrrole. Examples of ketones include, but are not limited to, acetone, 3-pentanone, dibenzyl acetone, 5-nonone, 4-heptanone, cyclohexanone, or cyclopentanone. Examples of Brønsted acids have been described herein.

An exemplary method is described in Example 3.

Lewis acid condensation to form calix[n]pyrrole where n is 4, 5, 6, 7, or 8.

A pyrrole and a ketone are reacted in the presence of a Lewis acid to form a calix[n]pyrrole where n is 4, 5, 6, 7, or 8. Examples of ketones and pyrroles have been described herein. Examples of Lewis acids include, but are not limited to, electron pair acceptors containing empty orbitals such as aluminum trichloride, boron trifluoride, gallium trichloride, or high valent metals such as titanium (IV), vanadium (V), tin (IV), manganese (III), or UO_2^{+2} . Exemplary methods include the methods described in Example 5.

Heterogeneous condensation to form calix[n]pyrroles where n is 4, 5, 6, 7 or 8.

A synthetic method for heterogeneous condensation to form calix[n]pyrroles where n is 4, 5, 6, 7 or 8 comprises reacting a pyrrole and a ketone in the presence of a heterogeneous catalyst to form a calix[n]pyrrole where n is 4, 5, 6, 7, or 8. Examples of pyrroles and ketone have been described herein. Examples of heterogeneous catalysts include, but are not limited to montmorillonite, zeolites, silica, alumina or a cation-exchange solid support. Exemplary methods include those methods found in Example 5.

Formation of pyrrole alcohols to form calix[n]pyrroles where n is 4, 5, 6, 7 or 8.

A method of synthesis of calix[n]pyrroles using pyrrole alcohols comprises reacting a ketone functionalized pyrrole with an alkyl metal or aryl metal to form a pyrrole alcohol, condensing the pyrrole alcohol with a Lewis acid, a Brønsted acid or a heterogeneous catalyst to form a cyclizable oligomer. Repeating the reaction of the ketone and the subsequent condensation n - 2 times to form a calix[n]pyrrole. Examples of pyrrole ketones include, but are not limited to, 2-acetyl-3,4-diethylpyrrole, 2-acetylpyrrole, 2-acetyl-3,4-dimethylpyrrole, 2-alkylacetyl-3,4-disubstituted pyrrole, 2-alkylacetyl-3-substituted pyrrole, or 2-alkylacetyl-4-substituted pyrrole. Examples of alkyl metals include, but are not limited to, alkyllithium, Grignard reagent, organocuprates, organozincs or organotin compounds. Examples of aryl metals include, but are not limited to, phenyllithium.

Using 2-acetylpyrrole and an alkyl or aryl lithium to illustrate the method described herein, 2-acetylpyrrole is reacted with butyl lithium to produce 1-butyl-1-methyl-1-(2-pyrrolo) methanol. 1-Butyl-1-methyl-1-(2-pyrrolo) methanol is condensed with 2-acetylpyrrole in the presence of a Brønsted acid. The reaction is repeated twice to form the calix[4]pyrrole.

Template synthesis

This synthetic method embodiment comprises reacting a ketone and pyrrole in the presence of a Lewis acid, a Brønsted acid or a heterogeneous catalyst in the presence of a

Me₃SiI. Non-covalent interactions may also be employed, such as carboxylic acid dimerization or metal complexation between the template and the ketone.

In the synthesis, the pyrrole condensation must be acid catalysed; the acid may be selected from Brønsted acids, for example, methanesulfonic acid, HCl, HClO₄, H₂SO₄, HNO₃, TFA, HCO₂H, or H₃PO₄ or Lewis acids such as VOCl₃, Mn(OAc)₃, SnCl₄, AlCl₃, or UO₂(OAc)₂. The synthesis is carried out in an organic solvent that is compatible with the particular ketone-containing template utilized. Solvents such as chloroform, dichloromethane, carbontetrachloride, toluene, benzene, and mixtures of these solvents with methanol, ethanol, propanol and the like may be used. The linker to the ketone could also be changed to include any spacer such as benzyl groups, alkynes, alkenes, ethers, polyethers, thioethers, bipyridyl, phenanthroline, anthracene, naphthalene, any polyaromatic, quinone, ferrocene, cobaltocenium or other metallocene, and the like.

Using calixarenes to illustrate the method of the present invention, the calix[4]arene skeleton is used as a template, around which a calix[4]pyrrole forms. Although calixarene-linked and calixarene-capped porphyrins as well as calixarenes linked to pyrrole groups are known, to the best of the present inventors knowledge, compound 15 represents the first example of a calixarene-capped calixpyrrole.

Calix[n]pyrroles derivatives, multimers, and conjugates

With use of either the *meso*-“monohook” acid 4 or the β-“monohook” acid 12 there exist possibilities of substitution through various functional groups. Formation of an amide bond, for example, could lead to attachment of natural nucleotides as well as the “unnatural” nucleotide analogues. Using nucleotide-functionalized calixpyrroles, it may be possible to do through-membrane transport of nucleotide analog drugs such as AZT, acyclovir and the like (see below). In addition to one “hook”, one may add two, three, four, five, six, seven or eight “hooks” at the *meso*- or β- positions of the calixpyrrole. Arrays of calixpyrroles spanning membranes and acting as anion channels analogous to cation channels are contemplated by the present inventors. By using poly-amine-containing compounds, dimers, trimers and tetramers of calixpyrrole could be used as carriers for transporting polyanionic species through aliphatic membranes. Through NMR work and work with HPLC, it has been determined that calixpyrroles show a higher affinity for phosphate anions than other oxyanions. With this in mind, arrays of calixpyrroles could be used to encircle RNA and DNA molecules, thus providing anti-viral activity. Specificity of binding could be gained by attaching specific nucleotides to the calixpyrrole oligomer. Attachment of known nucleotide cleaving agents such as Fe-EDTA could facilitate the site directed cleavage of RNA. Calixpyrrole also binds to carboxylic acid groups by HPLC and could be used for extraction of pesticide and herbicide metabolites. Additionally, with the addition of a cation-binding functionality, amino acid recognition and transport are possible. A solid silica-supported calixpyrrole has shown selectivity of various N-benzyl protected amino acids. This could lead to specific binding of

proteins and interruption of enzymatic pathways. The attachment of sugars, and mono- and polysaccharides to the calixpyrrole could allow for recognition of compounds such as glucose-6-phosphate.

By attachment of a redox-active reporter group; calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine macrocycles may function as potentiometric or amperometric sensors for cations, anions, or neutral molecules.

By attachment of a fluorophoric reporter group; calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine macrocycles may function as fluorescent sensors for cations, anions, or neutral molecules.

By attachment of a chromophoric reporter group; calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine macrocycles may function as chromophoric sensors for cations, anions, or neutral molecules.

Calix[n]pyrroles, calix[m]pyridino[n]pyrroles and calix[m]pyridines as separation media

Separation of closely similar molecules from highly complex mixtures requires high resolution techniques. Ion exchange, affinity chromatography, reverse phase chromatography, and gel filtration represent current types of separation techniques. The present macrocycles provide a new type of separation media due to non-covalent, primarily hydrogen bonding, interaction between the macrocycle and a guest molecule.

In ion exchange, separation is based on the charges carried by the solute molecules. An ion exchanger includes an insoluble matrix or solid support to which charged groups have been covalently bound. The charged groups are associated with mobile counterions. The counterions can be reversibly exchanged with other ions of the same charge without altering the matrix. It is possible to have both positively and negatively charged exchangers. Positively charged exchangers have negatively charged ion (anions) available for exchange and are so termed cation exchangers. Negatively charged exchangers have positively charged counterions and are termed anion exchangers. The matrix may be based on inorganic compounds, synthetic resins, polysaccharides, silicas, aluminas, or polymeric supports. The nature of the matrix determines its physical properties, such as its mechanical strength and flow characteristics, its behavior towards test samples, and to some extent, its capacity.

Affinity chromatography is a type of adsorption chromatography in which the molecule to be purified is specifically and reversibly adsorbed by a complementary binding substance (ligand) immobilized on an insoluble support (matrix). Cyanogen bromide activation has been the most common method for coupling ligands to a matrix. A successful separation requires that a ligand is available and that it can be covalently attached to a chromatographic bed material. It is also important that the immobilized ligand retains its specific binding affinity for the substance of interest and that methods are available for selectivity desorbing the bound substances in an active form, after washing away unbound material.

Reverse phase chromatography relies on hydrophobic surfaces, especially silylated alkyl (R₁, R₂, R₃, SiO), on the solid support to bind hydrophobic molecules. More polar molecules pass through the column unimpeded. The hydrophobic media may include aromatic molecules, long chain aliphatic hydrocarbons, or cyclic hydrocarbons.

5 Separation in gel filtration depends on the different abilities of the various sample molecules to enter pores that contain a stationary phase. As a solute passes down a chromatographic bed, its movement depends upon the bulk flow of the mobile phase and upon the Brownian motion of the solute molecules which causes their diffusion into and out of the stationary phase. Very large molecules that never enter the stationary phase move through the chromatographic bed fastest. Smaller molecules that can enter the gel pores move more slowly through the column since they spend a proportion of their time within the stationary phase. Molecules are, therefore, eluted in the order of decreasing molecular size.

In contrast to the known methods of chromatography, the macrocycles of the present invention provide new principles for separation of ions and molecules. Interactions between macrocycles of the present invention and ions and molecules are noncovalent, primarily hydrogen bonding. As a result, surprising and unexpected affinities and selectivities are observed for the separation of very similar molecules. For instance, the following sets of anions or neutral molecules were successfully separated (Example 10):

- Phenyl arsenate, benzene sulfonate, benzoate, and phenyl phosphate;
- 20 • 5'-Adenosine monophosphate, 5'-adenosine diphosphate and 5'-adenosine triphosphate;
- Uridine, phenol, aniline, benzene, and 4 iodo-nitro benzene;
- Cbz-N-protected-serine, glutamine, alanine, phenylalanine, tryptophan, aspartate, and glutamate;
- 4-Fluorobiphenyl, 4,4'-difluorobiphenyl, 2,2',3,3',5,5',6,6',-octafluorobiphenyl, and perfluorobiphenyl;
- 25 • A mixture of polydeoxythymidylates oligomers containing between (12) and (18) bases;
- A mixture of polydeoxythymidylates oligomers containing between (19) and (24) bases;
- TCTAGA, GCATGC, and CCCGGG;
- Phthallate, isophthallate and teraphthallate;
- 30 • Benzoate, isophthallate, and 1,3,5-tricarboxybenzene; and
- Carbofuran, carbendazim, bromacil, bentazon, carboxin, and norflurazon.

A method of use of the present invention comprises separating a first molecule, anion or cation from a mixture of a first molecule, anion or cation and other species using a form of solid support-immobilized calix[n]pyrrole, calix[m]pyridino[n]pyrrole or calix[m]pyridine by contacting the solid support with the mixture to separate a first molecule, anion or cation from a mixture of a first molecule, anion or cation and other species. Examples of solid supports include, but are not limited to, silica gel, aminopropyl silica gel, alumina, polyacrylamide polymer beads, polystyrene polymer beads, sepharose, sephadex, agarose, clays, or zeolites

coupled to calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines. Examples of forms of solid supports include, but are not limited to chromatography columns, thin-layer chromatographic supports, electrophoresis gels, or capillary electrophoresis tubes. Examples of ions or neutral molecules to be separated are neutral aromatic or aliphatic species (polymeric, oligomeric and monomeric), nucleotides, oligonucleotides, polyoxometallates, or anionic species (polymeric, oligomeric and monomeric) such as pertechnate, a polyhalobiphenyl (including polychlorobiphenyl) or inorganic phosphate.

The inventors' discovery that calix[n]pyrroles recognize and bind phosphates led the inventors to reason that such macrocycles would be ideal for use in laboratory techniques to separate and purify oligonucleotides.

The present inventors realized that for a solid support to be most useful, it must not only separate similar nucleotides and oligonucleotides from one another, it must do so rigorously and correctly based on chain length or size. Unfortunately, prior to the present invention, there were no suitable phosphate chelating agents which made this an achievable goal for use in either HPLC or electrophoretic separation systems. As described in the examples herein, the inventors designed and constructed calixpyrrole-based supports and used them to achieve the separation of nucleotides and oligonucleotides, thus overcoming many of the existing drawbacks.

A further utility of such calixpyrrole-substituted materials is their use as tools in the removal of phosphorylated environmental contaminants from ground water, soil, foodstuffs, and the like. They may therefore be employed to analyze and separate pesticides such as but not limited to carbofuran, carbendazim, bromacil, bentazon, carboxin and norflurazon. Calixpyrrole-substituted silica gels and columns may even be employed in the rapid detection and analysis of organophosphorus chemical warfare agents, allowing them to be disposed of where necessary.

Another technical area in which significant improvements could be made is in oligonucleotide analysis. For example, automated gene sequencing is currently carried out using either radio- or fluorescent labeled nucleic acid gel electrophoresis. This technique is limited by the requirement for either slab or tubular polyacrylamide gels. An alternative approach, currently being considered on a research basis, is to use capillary electrophoresis. Unfortunately, this is limited when it comes to achieving separations based purely on electrostatics and oligomer size. A technique which would allow for such separation would therefore represent a significant improvement in this area.

Two new derivatized silica gels have been produced. Gel **M** is linked to the appended calixpyrroles *via* a calixpyrrole *meso* carbon and Gel **B** is linked to the appended calixpyrrole *via* a calixpyrrole β -carbon. Both Gels have been demonstrated to separate mixtures of anions such as oligonucleotides (dT₁₂₋₂₄), nucleotides, aromatic polycarboxylates, phenyloxanions, halides, phosphates and sulfates. Additionally, the calixpyrrole-modified silica gels can separate

mixtures of neutral molecules such as polyfluorobiphenyls and simple aromatic species. Thus the calixpyrrole modified column supports offer a wide range of separatory ability.

The calix[n]pyrrole may be associated with a metal that is bound by substituents on the periphery of the macrocycle. Heterocyclic macrocycles containing free electron pairs chelate cations. Pyridine is known to coordinate to certain metal cations. Calix[m]pyridines also contain nitrogen atoms that are expected to donate electron density to cations. This donation would require that the cations stay in close proximity to the pyridine units, thereby, binding the cations to the core of the ring.

In vivo applications

The present inventors envision use of calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines in DNA recognition and modification. Water-soluble calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines may be particularly advantageous for use in a number of ways, such as in cellular recognition, targeting, and in the transport of biologically important molecules. Generally, water-soluble calix[n]pyrroles are preferred for biomedical applications. "Water soluble" means soluble in aqueous fluids to about 1 mM or better.

Anionic phosphorylated entities are ubiquitous in biology. They play a critical role in a variety of fundamental processes ranging from gene replication to energy transduction. In addition, certain phosphate-bearing nucleotide analogues, such as, e.g., 9-(β -D-xylofuranosyl)guanine-5'-monophosphate (xylo-GMP), are known to display antiviral activity *in vitro*. However, xylo-GMP, like a considerable number of phosphorylated nucleotide analogues which exhibit anti-viral activity in cell-free extracts, is inactive *in vivo* due to its inability to cross lipophilic cell membranes.

The anti-herpetic agent, acyclovir (9-[(2-hydroxyethoxy)methyl]-9H-guanine), is active *in vivo*. Acyclovir can enter the cell only in its uncharged nucleoside like form. Once in the cytoplasm, it is phosphorylated, first by a viral encoded enzyme, thymidine kinase, and then by relatively non-specific cellular enzymes to produce an active ionic triphosphate nucleotide-like species. There, it functions both as an inhibitor of the viral DNA polymerase and as a chain terminator for newly synthesized herpes simplex DNA.

The biological limitations of many other potential antiviral agents, including xylo-G, arise from the fact they are not phosphorylated once inside the cell and are therefore largely or completely inactive. If, however, the active monophosphorylated forms of these putative drugs could be transported into cells, it would be possible to fight viral infections with a large battery of otherwise inactive materials. If such specific into-cell transport were to be achieved, it would therefore greatly augment the treatment of such debilitating diseases as, for example, AIDS, herpes, hepatitis and measles. Given the fact that AIDS is currently a major world health problem of frightening proportions, and that something so nominally benign as measles still claims over 100,000 lives per year world-wide, treatment of these diseases would be particularly timely and worthwhile.

A major aim of the inventors' studies has been therefore to provide a means of transporting active mono- and phosphorylated forms of these and other agents into cells. This would allow a wide range of otherwise inactive compounds, such as antivirals, to be employed therapeutically, and would also create new possibilities for gene therapy.

5 Calix[n]pyrrole- or calix[m]pyridino[n]pyrrole-based systems may be made effective as neutral regime carriers, say, e.g., for GMP by constructing a of polytopic receptor system in which a nucleobase recognition unit, in this case, a cytosine-like group, is appended directly to a phosphate-chelating calix[n]pyrrole or calix[m]pyridino[n]pyrrole macrocycle. Nucleobase recognition units are demonstrated herein for use in the specific binding and transport of
10 complementary nucleobases and nucleobase-containing compounds.

To synthesize multitopic receptors, calix[n]pyrrole conjugates were developed to address the following objectives: (i) the independent development of molecular recognition strategies for the complexation of two very different kinds of substrates (charged anionic and neutral nucleobase); (ii) subsequent co-combination so as to provide receptors bearing both
15 kinds of binding subunits; and (iii) various alternative methods of receptor oligomerization so as to provide oligomeric species bearing numerous combinations of multitopic receptors.

Pursuing these strategies led to the development of the calix[n]pyrrole-based ditopic receptor systems of the present invention, capable of recognizing both the anionic phosphate and the neutral portions of the nucleotide derivatives, such as the purine or pyrimidine moieties.
20 The present inventors envisage expanding this theme to the preparation of oligomeric, multitopic, receptors capable of recognising multiple phosphate anions and nucleobase portions of nucleotide derivatives arranged in specific sequences.

Ditopic receptor systems are ideal vehicles for the intracellular transport of oligonucleotides and their derivatives, including anti-viral agents. The multitopic receptors,
25 likewise, are contemplated to be of use in binding to oligonucleotides and specific sections of DNA or RNA and in transporting such nucleic acid segments into cells.

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in
30 the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Pharmaceutical preparations

35 For *in vivo* and *ex vivo* uses, calix[n]pyrroles, calix[m]pyridino[n]pyrroles and calix[m]pyridines are provided as pharmaceutical preparations. A pharmaceutical preparation of a calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines may be administered alone or in combination with pharmaceutically acceptable carriers, in either single or multiple

doses. Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. The pharmaceutical compositions formed by combining a calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines of the present invention and the pharmaceutically acceptable carriers are then easily administered in a variety of dosage forms. Administration may be intravenous, intraperitoneal, parenteral, intramuscular, subcutaneous, oral, or topical, with intravenous administration being preferred.

Solutions of the calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines in sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solution may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. Topical creams, emulsions, solutions, and the like are contemplated for applications to surface areas of the body. Topical application may also be by iontophoresis.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy use with a syringe exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars such as mannitol or dextrose or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder

of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, permeation enhancers, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

Calix[n]pyrroles, calix[m]pyridino[n]pyrroles or calix[m]pyridines may be incorporated into liposomes for use in the present invention. Liposomes may be prepared by any number of techniques that include freeze-thaw, sonication, chelate dialysis, homogenization, solvent infusion, microemulsification, spontaneous formation, solvent vaporization, reverse phase, French pressure cell technique, or controlled detergent dialysis, for example. Preparation may be carried out in a solution, such as a phosphate buffer solution, containing calix[n]pyrrole-, calix[m]pyridino[n]pyrrole or calix[m]pyridine-lipophilic molecule conjugates so that the conjugate is incorporated into the liposome membrane. Alternatively, the conjugate may be added in already formed liposomes. Liposomes employed in the present invention may be of any one of a variety of sizes, preferably the less than about 100 nm in outside diameter, more preferably less than about 50 nm.

Micelles may be prepared by suspension of a calix[n]pyrrole-, calix[m]pyridino[n]pyrrole- or calix[m]pyridine-lipophilic molecule and lipid compound(s) in an organic solvent, evaporation of the solvent, resuspension in an aqueous medium, sonication and then centrifugation. Alternatively, the calix[n]pyrrole-, calix[m]pyridino[n]pyrrole- or calix[m]pyridine-lipophilic molecule may be added to preformed micelles.

Techniques and lipids for preparing liposomes and micelles are discussed in U.S. Patent 5,466,438, incorporated herein by reference.

Use of calix[n]pyrroles, calix[m]pyridino[n]pyrroles, or calix[m]pyridines in ex vivo and in vivo treatment of body tissues.

The method of invention comprises the dialysis bodily fluids using calix[n]pyrroles, calix[m]pyridino[n]pyrroles, or calix[m]pyridines. Examples of dialysible substrates include, but are not limited to, urea, phosphate containing molecules or halide waste. Examples of conditions for treatment include gout, diabetes or drug overdoses.

Using kidney dialysis to illustrate, but not limit, the methods described above, bodily fluid would be passed through a membrane containing calix[n]pyrroles, calix[m]pyridino[n]pyrroles, or calix[m]pyridines. The relevant wastes such as, but not limited to, inorganic phosphate, would then be removed.

Example 1
Novel Crystal Structures of Complexed Prior-Art
***meso*-Substituted Calixpyrrole Macrocycles**


5 The present inventors have crystallized anion- and neutral molecule-complexed forms of known calix[4]pyrrole macrocycles and have studied the coordination properties of calixpyrroles in solution and in the solid state. As a result, the present inventors have demonstrated unexpected crystalline forms of calixpyrrole macrocycles and the existence of a conformational change between an unbound macrocycle and a macrocycle bound to an anion or a
10 neutral substrate. In the solid state, uncomplexed calix[4]pyrrole macrocycles adopt an alternate conformation where adjacent pyrrole rings are oriented in opposite directions. Solid-state complexed calix[4]pyrrole macrocycles are shown herein to adopt a cone conformation such that the four NH protons of the pyrrole rings can hydrogen bond to the bound substrate. In solution, a dynamic equilibrium exists where the bound anionic or neutral substrate is
15 reversibly bound to all or less than all of the pyrrole NH groups. Such a conformational change would alter the properties of the calixpyrrole, for example, groups on the *meso*- or β -positions would be brought into closer proximity in the cone configuration. This conformational change may be taken advantage of in the design of a sensor, for example.

meso-Octamethylcalix[4]pyrrole **1** was prepared using prior art procedures in the
20 following manner. Pyrrole (20 g) and acetone (20 g) were introduced into a stirred 500 mL round-bottomed flask fitted with a reflux condenser. Methanesulfonic acid (1 mL) was added down the reflux condenser, causing a violent reaction. CARE! - The acid must be added slowly! Once the acid was added, the solid mass formed was broken up and a further 20 mL of acetone was added. The mixture was then heated at reflux for 30 min. After this time, the
25 mixture was allowed to cool and the white precipitate was filtered off.


 The white precipitate was crystallized from acetone to give a crude crystalline material. This was recrystallized from acetone, giving crystals of the product octamethylcalix[4]pyrrole **1**, suitable for single crystal X-ray diffraction studies.

meso-Tetraspirocyclohexylcalix[4]pyrrole **2** was prepared according to prior art
30 procedures as follows. A solution of 16.2 g (0.242 mol) of pyrrole and 20 mL of absolute ethanol was placed in three-necked 500 mL flask. The flask, fitted with a reflux condenser, was placed in an ice bath. To the stirred solution was added dropwise 1 mL of 37% aqueous hydrochloric acid solution, followed by 23.7 g (0.242 mol) of cyclohexanone over a period of 20 min. The stirrer was stopped by the solid formed in the flask when the addition of
35 cyclohexanone was complete. After standing at room temperature for 1 hour, the product was washed with 50 mL of cold water. The solid was removed from the flask by partial dissolution in ether. The total yield of the crude synthesis was 12.86 g (9.0%).

 Several crystallizations of the product from benzene and then from hexane gave white crystals of tetraspirocyclohexylcalix[4]pyrrole **2**, m.p. 272.0 - 272.5°.



1



2

X-ray Crystallographic Data. Single crystals suitable for X-ray crystallographic analysis were obtained by slow evaporation of acetone and acetone/dichloromethane (1:1 v/v) solutions of calixpyrroles 1 and 2, respectively. The crystallographic summary for 1 is as follows. Pale yellow crystals, tetragonal, $P4_1$, $Z = 2$ in a cell of dimensions $a = 10.097(1)$ Å, $c = 23.764(1)$ Å, $V = 2422.7(4)$ Å³, $\rho_{\text{calc}} = 1.18$ g cm⁻³, $F(000) = 928$. Data were collected at 198K on a Siemens P4 diffractometer, equipped with a Nicolet LT-2 low-temperature device and using graphite monochromatized Mo $K\alpha$ radiation ($\lambda = 0.71073$ Å). Data were collected using ω scans with a scan range of 1° in ω . 3733 unique reflections, 2552 with $F_o > 4(\sigma(F_o))$. The final $R = 0.0556$, $R_w = 0.132$, goodness of fit = 0.999 for 290 parameters. The crystallographic summary for 2 is as follows (CH_2Cl_2). Colorless crystals, monoclinic, $P2_1/c$, $Z = 4$ in a cell of dimensions $a = 10.993(2)$ Å, $b = 14.250(3)$ Å, $c = 23.466(4)$ Å, $\beta = 97.254(14)^\circ$, $V = 3646.5(11)$ Å³, $\rho_{\text{calc}} = 1.23$ g cm⁻³, $F(000) = 1448$. Data were collected on a Siemens P3 diffractometer. 5562 unique reflections, 2285 with $F_o > 4(\sigma(F_o))$. The final $R = 0.0762$, $R_w = 0.141$, goodness of fit = 0.995 for 424 parameters. As illustrated in FIG. 1A and FIG. 1B, molecules 1 and 2 adopt a 1,3-alternate conformation in the solid state wherein adjacent rings are oriented in opposite directions similar to a previously-reported octaethyl analogue.

Crystals of anion complexes of the calixpyrroles 1 and 2 were then prepared in order to study the conformational changes from the free ligand to the complexed ligand. Crystals of the tetrabutylammonium chloride complex of calixpyrrole 1 were obtained by slow evaporation of a dichloromethane solution containing an excess of the chloride salt. The crystallographic summary for $1 \cdot (\text{nBu}_4\text{NCl}) \cdot \text{CH}_2\text{Cl}_2$ is as follows: Colorless crystals, Orthorhombic, $Pna2_1$, $Z = 4$ in a cell of dimensions $a = 21.247(3)$ Å, $b = 19.937(2)$ Å, $c = 10.691(1)$ Å, $V = 4528.7(9)$ Å³, $\rho_{\text{calc}} = 1.16$ g cm⁻³, $F(000) = 1720$. 6027 unique reflections, 3566 with $F_o > 4(\sigma(F_o))$. The final $R = 0.0959$, $R_w = 0.246$, goodness of fit = 1.224 for 478 parameters. Crystals of the tetrabutylammonium fluoride complex of compound 2 were obtained using a similar procedure and the crystallographic summary for $2 \cdot (\text{nBu}_4\text{NF})$ is as follows. Colorless crystals, tetragonal, $P4/nnc$, $Z = 4$ in a cell of dimensions $a = 15.732(1)$ Å, $c = 20.076(1)$ Å, $V = 4968.7(5)$ Å³, $\rho_{\text{calc}} = 1.14$ g cm⁻³, $F(000) = 1872$. 3634 unique reflections, 2220 with $F_o > 4(\sigma(F_o))$. The final $R = 0.064$, $R_w = 0.171$, goodness of fit = 1.029 for 152 parameters.

The structures so obtained (FIG. 2A and FIG. 2B) reveal that in both cases the calix[4]pyrrole ligand adopts a cone-like conformation such that the four NH protons can hydrogen bond to the bound halide anion. While these two structures are thus grossly similar, in the case of the chloride complex (FIG. 2A) the nitrogen-to-anion distances are in the range of 3.264(7) - 3.331(7) Å, while for the corresponding fluoride complex they are 2.790(2) Å (FIG. 2B) (the four pyrrole groups are equivalent by symmetry). As a result, in these two complexes the chloride and fluoride anions reside 2.319(3) Å and 1.499(3) Å above the N_4 root mean square planes of calixpyrroles **1** and **2**, respectively. Thus, fluoride anion appears to be more tightly bound in the solid state.

Expanding the range of potential guests, complexes of neutral species were then prepared. Crystal complexes of calixpyrrole **1** and *N,N*-dimethylformamide ("1•2(DMF)") were grown by dissolution of compound **1** in DMF and subsequent crystallization. The $N_{\text{pyrrole}} \cdots O_{\text{DMF}}$ distances are 2.908(2) and 2.924(2) Å, with associated N-H...O angles of 167(2)° and 166(2)°, respectively. The unsaturated portion of the amide lies 3.363(3) Å above the plane of a third pyrrole ring (the planar twist angle between these two moieties is 7.1(1)°, suggesting that a π -stacking interaction helps to stabilize 1•2(DMF). The crystallographic summary for 1•2(DMF) is as follows: colorless prisms from DMF, triclinic space group $P\bar{1}$, $Z = 1$, $a = 9.473(1)$ Å, $b = 9.573(1)$ Å, $c = 10.316(1)$ Å, $\alpha = 66.428(4)^\circ$, $\beta = 80.987(5)^\circ$, $\gamma = 72.025(4)^\circ$, $V = 815.00(14)$ Å³, $\rho_{\text{calc}} = 1.17$ g cm⁻³, $F(000) = 312$. Final $R = 0.0390$, $R_w = 0.0952$, GOF = 1.061 for 291 parameters. The NH protons of 1•2(DMF) were obtained from a ΔF map and refined with isotropic thermal parameters.

Crystal complexes of calixpyrrole **1** and methanol ("1•2(CH₃OH)") have also been studied by X-ray crystallography. The crystals were prepared by slow evaporation of a methanol/dichloromethane solution of the ligand. The calixpyrrole unit in 1•2(MeOH) was found to assume a 1,3-alternate conformation in the solid state. Single molecules of the alcohol lie above and below the macrocycle, each one held in place by hydrogen bonds to two pyrrolic NH groups. The four symmetry-equivalent H-bonds ($N_{\text{pyrrole}} \cdots O_{\text{MeOH}}$) are 3.155(4) Å long, a value which is very close to the $N_{\text{pyrrole}} \cdots O_{\text{MeOH}}$ distances (*ca.* 3.0-3.2 Å) found in a methanol complex of a tetrapyrrolic "expanded porphyrin." Further evidence that methanol is bound to **1**, rather than merely occupying space in the lattice, is given by the inward tilt of the pyrroles of 1•2(MeOH). This effect compresses the "cross ring" $N_1 \cdots N_{1C}$ and $N_{1A} \cdots N_{1B}$ distances by *ca.* 0.15 Å relative to those in free **1**, although it is not sufficient to allow for a linear alignment of the three hydrogen bonding atoms ($N_{\text{pyrrole}}\text{-H} \cdots O_{\text{MeOH}} = 152.1(4)^\circ$). Crystal data for 1•2(MeOH): yellow prisms from CH₂Cl₂-MeOH, tetragonal space group $I\bar{4}$, $Z = 2$, $a = 10.383(2)$ Å, $b = 10.383(2)$ Å, $c = 13.232(5)$ Å, $V = 1426.6(6)$ Å³, $\rho_{\text{calc}} = 1.15$ g cm⁻³, $F(000) = 536$. Final $R = 0.0787$, $R_w = 0.205$, GOF = 1.099 for 84 parameters. The NH proton of 1•2(MeOH) was calculated in an idealized position (N-H = 0.90 Å) with U_{iso} set to 1.2 U_{eq} for the attached N atom.

Crystal complexes of calixpyrrole **1** and dimethylsulfoxide ("1•(DMSO)") have also been studied by X-ray crystallography. The crystals also undergo a phase change as they are cooled, at room temperature there is one crystallographically distinct calixpyrrole in the unit cell, at -75 °C there are two. In both phases the calixpyrrole adopts a 1,3-alternate conformation wherein a DMSO molecule is hydrogen bonded to one pyrrole NH group (at low temperature N...O = 2.962, 2.985 Å; H...O = 2.089, 2.024 Å and N-H...O = 176.2° and 174.5° and at room temperature N...O = 2.996 Å, H...O = 2.099 Å and N-H...O = 174.5°). Crystallographic data for 1•(DMSO): There are two crystallographic modifications of 1•(DMSO). The room temperature modification is monoclinic, $P2_1/c$, $Z = 4$, in a cell of dimensions: $a = 10.423(2)$, $b = 23.599(6)$, $c = 12.480(2)$ Å, $\beta = 107.72(1)^\circ$, $V = 2924(1)$ Å³, $\rho_{\text{calc}} = 1.15$ g cm⁻³, $F(000) = 1096$, 6720 unique reflections, 2161 with $F_o > 4(\sigma(F_o))$. The final $R(F) = 0.0866$, $R_w(F^2) = 0.253$, goodness of fit = 0.981 for 335 parameters refined on F^2 . The low-temperature modification (-75 °C) is triclinic, $P\bar{1}$, $Z = 4$, in a cell of dimensions: $a = 10.362(3)$, $b = 12.373(7)$, $c = 23.468(10)$ Å, $\alpha = 89.77(4)$, $\beta = 88.35(3)$, $\gamma = 72.36(4)^\circ$, $V = 2866(2)$ Å³, $\rho_{\text{calc}} = 1.17$ g cm⁻³, $F(000) = 1096$, 10061 unique reflections, 5213 with $F_o > 4(\sigma(F_o))$. The final $R(F) = 0.0819$, $R_w(F^2) = 0.212$, goodness of fit = 1.075 for 700 parameters refined on F^2 .

The complex of calix[4]pyrrole **1** and *N,N*-dimethylformamide ("1•2(DMF)") indicates that individual symmetry-equivalent guests are found above and below the host, and that each amide is hydrogen bonded to adjacent pyrroles (FIG. 3A). The conformation of the calixpyrrole is therefore 1,2-alternate. To the present inventors' knowledge, 1•2(DMF) is the first calixpyrrole in this conformation to be unambiguously characterized. The complexes of 1•2(CH₃OH) (FIG. 3B) and 1•(DMSO) (FIG. 3C) both adopt the 1,3-alternate conformation in the solid state. However in the case of 1•2(CH₃OH) all the pyrrole NH groups are involved in hydrogen bonding, whereas in the case of 1•(DMSO) only one pyrrolic NH group hydrogen bonds to the calixpyrrole.

While cone, 1,3-alternate, 1,2-alternate and partial cone conformations for calixpyrrole macrocycles have been proposed previously, the present studies provide the first demonstration of the crystalline forms of anion-complexed molecules. Solution phase data are provided in Example 10.

Example 2


Novel *meso*-Substituted Calix[4]pyrrole Macrocycles

The present example provides *meso*-substituted calix[4]pyrrole macrocycles that have not previously been known. In particular, calix[4]pyrrole macrocycles having functional groups for further derivatization have proved particularly useful as provided for in later examples.

Calix[4]pyrrole *meso*-monoester 3. 4-Acetylbutyric acid (3.8 g, 29 mmol) was dissolved in methanol (50 mL) in a 100 mL round bottomed flask with magnetic stirring. Concentrated hydrochloric acid (1 mL, 37%) was added and the mixture heated at reflux for 12

hr. The solvent was then removed *in vacuo* and the residue redissolved in dichloromethane (50 mL). The solution was then washed with saturated NaHCO_3 (aq) solution (2 x 50 mL) and then brine (1 x 100 mL). The organic layer was separated and then dried over MgSO_4 and reduced *in vacuo* to give the methyl ester, methyl-4-acetylbutyrate, as a yellow oil (3.5g, 24.3 mmol) corresponding to a yield of 84%. ^1H NMR (CDCl_3) 3.64 (s, 3, OCH_3), 2.49 (t, $J = 7.2$ Hz, 2, CH_2), 2.32 (t, $J = 7.2$ Hz, 2, CH_2), 2.12 (s, 3, CH_3), 1.86 (m, 2, CH_2). ^{13}C NMR (CDCl_3) 207.0, 172.8, 50.7, 41.7, 32.3, 29.1, 18.25. CI MS m/z @ 145 (MH^+)


Methyl-4-acetylbutyrate (0.80 g, 5.6 mmol) was dissolved in methanol (50 mL) in a 100 mL round bottomed flask. Cyclohexanone (0.55 g, 5.6 mmol) and pyrrole (0.75 g, 11.1 mmol) were added and the mixture cooled in an ice bath to 0°C . The flask was fitted with a reflux condenser and magnetically stirred. Methanesulfonic acid (20 drops) was added dropwise over the course of five minutes, and the mixture was stirred at 0°C for 1 hr. and then at room temperature overnight. The solvent was removed *in vacuo*. FAB mass spectroscopic analysis showed the presence of mono-, di-, tri- and tetrahook material in the crude reaction mixture as well as tetraspirocyclohexylcalix[4]pyrrole. The residue was purified by column chromatography on silica gel using dichloromethane as eluant. This yielded the monohook calixpyrrole methyl ester **3** as a white powder (211mg, 0.33 mmol) corresponding to a yield of 11.8%. ^1H NMR (CDCl_3) 7.07 (br. m, 4, NH), 5.88 (m, 8, $\text{CH}(\text{py})$), 3.63 (s, 3, OCH_3), 2.22 (t, $J = 7.2$ Hz, 2, CH_2), 1.93 (v. br., 14, $6\text{CH}_2(\text{cyclohexyl}) + \text{CH}_2(\text{hook})$) 1.56-1.35 (v. br., 23, $\text{CH}_2(\text{hook}) + 9\text{CH}_2(\text{cyclohexyl}) + \text{CCH}_3$). ^{13}C NMR (CDCl_3) 173.8, 137.169, 136.5, 136.3, 130.7, 103.6, 103.5, 103.2, 53.4, 51.37, 39.7, 39.5, 39.4, 38.5, 37.5, 37.2, 36.9, 36.8, 34.0, 26.1, 25.9, 22.75, 22.70, 19.9. CI MS m/z @ 635 (MH^+). High Resolution Mass Spectrum: Mass = 634.4236 mDa 1.1 ppm 1.8 Calc. mass 634.4247 DBE 17.0 $\text{C}_{41}\text{H}_{54}\text{N}_4\text{O}_2$.



Calix[4]pyrrole meso-monoacid 4.

The calixpyrrole methyl ester (211 mg, 0.33 mmol) (compound **3**) was dissolved in 20 mL ethanol in a 100 mL round bottomed flask with magnetic stirring. The flask was fitted with a reflux condenser and the solution heated to reflux. An aqueous solution of sodium hydroxide

(20 mL 2.0 M) was added and the mixture heated at reflux for 4 hr. The solution was then allowed to cool and the ethanol present in solution was removed *in vacuo*. The remaining aqueous solution was acidified to pH 1 with perchloric acid (70%). The product was then extracted from this aqueous phase with dichloromethane (50 mL). The dichloromethane layer was washed with water (100 mL) and dried over MgSO₄. Removal of the solvent afforded the “monohook” calixpyrrole acid **3** as an off-white powder (187 mg, 0.30 mmol) corresponding to a yield of 90.6%. ¹H NMR (CDCl₃) 7.02 (br. m., 4, NH), 5.89 (m, 8, CH_(py)), 2.26 (t, 2, CH₂), 1.91 (br., 14, 6CH₂(cyclohexyl) + CH₂(hook)), 1.44 (br. m., 24, 9CH₂(cyclohexyl) + CH₂(hook) + CCH₃) + 0.5H₂O) ¹³C NMR (CDCl₃) 179.4, 137.3, 136.6, 136.2, 103.7, 103.6, 103.5, 103.2, 53.4, 39.7, 39.4, 39.3, 38.5, 37.4, 37.2, 36.9, 36.8, 33.8, 26.0, 25.9, 22.7, 22.6, 19.5. FAB MS *m/z* @ 620 (MH⁺). High Resolution Mass Spectrum: Mass = 620.4073 mDa 1.7 ppm 2.8 Calc. mass 620.4090 DBE 17.0 C₄₀H₅₂N₄O₂.




X-ray quality crystals of the tetrabutylammonium salt of calix[4]pyrrole carboxylate **4** were obtained by slow evaporation of a dichloromethane solution of **4** in the presence of excess tetrabutylammonium fluoride trihydrate. The resulting crystal structure demonstrated that this anionic calix[4]pyrrole self-assembles into dimers in the solid state. The carboxylate moiety is hydrogen bonded to the pyrrole array of an adjacent calix[4]pyrrole (FIG. 4). Therefore the calix[4]pyrrole **4** adopts a cone conformation in the dimer. The separation between the root mean square planes of the nitrogen atoms in each macrocycle is 7.41(1)Å. Crystallographic summary for (C₄₀H₅₁N₄O₂)¹⁻ (C₁₆H₃₆N)¹⁺ · 2CH₂Cl₂, colorless needles were grown from CH₂Cl₂, triclinic, P $\bar{1}$, Z=2 in a cell of dimensions: *a* = 12.682(2), *b* = 13.127(2), *c* = 17.871(2)Å, α = 99.913(9), β = 90.527(9), γ = 98.49(1)°, *V* = 2896.7(6)Å³, ρ_{calc} = 1.18 g cm⁻³, F(000) = 1116. A total of 11364 reflections were measured, 10078 unique (*R*_{int} = 0.044) on a Siemens P4 diffractometer using graphite monochromatized Mo K α radiation (λ = 0.71073Å). The structure was refined on *F*² to an *R*_w = 0.220, with a conventional *R* = 0.084, with a goodness of fit = 1.031 for 661 refined parameters. The geometry of the interaction is: N1-H1N ... O1 (related by 1-x, 1-y, 1-z), N...O 2.975(5)Å, H...O 2.19(4)Å, N-H...O 162(4)°; N2-H2N ... O1, N...O 2.980(5)Å, H...O 2.18(5)Å, N-H...O 167(4)°; N3-H3N ... O1, N...O

2.950(5)Å, H...O 2.14(4)Å, N-H...O 163(4)°; N4-H4N ... O1, N...O 2.916(5)Å, H...O 2.09(4)Å, N-H...O 176(4)°.

"Monohook" acid **4** is particularly useful as a synthon for the production of calix[4]pyrrole derivatives, conjugates, modified silica gels and the like. Additionally the present inventors expect that compound **4** may be used as a zwitterion binding agent as cations may coordinate to the carboxylate functionality and anions to the calixpyrrole core.

Tetravinylcalix[4]pyrrole 5. Pyrrole (0.745 g, 11.12 mmol) and 5-hexan-2-one (1.09 g, 11.12 mmol) were dissolved in methanol (100 mL). Methanesulfonic acid (0.5 mL) was added and the reaction was stirred in the dark for 12 hr. The solvent was then removed *in vacuo* and the reaction mixture was purified by column chromatography on silica gel using dichloromethane as eluant. The first fraction from the column was collected and, upon removal of the solvent, afforded a colorless oil (1.05 g, 64% yield).


The product **5** is a mixture of conformers of tetravinylcalix[4]pyrrole where the groups in the *meso* positions are each in one of two orientations, with the bulkier alkene group either above or below the plane of the *meso*-carbon atoms (see Example 7). ¹H NMR (CDCl₃) : 7.02 (br. s, 4H, NH), 5.91 (m, 8H, CH_(pyrrole)), 5.71 (m, 4H, =CH-), 4.90 (m, 8H, =CH₂), 2.22 (br. m., 16H, CH₂CH₂). ¹³C NMR (CD₂Cl₂) : (139.4, 139.3, 139.2), (137.83, 137.80, 137.70, 137.63, 137.58), (114.5, 114.4, 114.35), (104.4, 104.3, 104.26, 104.2, 104.1, 104.0), (40.02, 39.98), (39.2, 39.0), 29.2, (26.45, 26.43, 26.23, 26.15, 26.04, 25.97).



Calix[4]pyrrole **5** may prove to be particularly useful as a monomer for use in the synthesis of calixpyrrole polymer materials. A known method uses pyrrole, a C₄-C₆ saturated alicyclic ketone and an acid containing vinyl groups or triple bonds to form a polymerized resin. In this case, the resulting products are undefined since it appears to be unknown where the modifying group is attached to the product. In contrast, the present method allows synthesis of monomer **5** which can then be subjected to polymerization. This method of forming a defined calix[n]pyrrole polymer is applicable for n = 4 - 8.

Meso-octabenzylcalix[4]pyrrole 6. Pyrrole (0.745 g, 11.12 mmol) and 1,3-diphenylacetone (2.34 g, 11.12 mmol) were dissolved in methanol (250 mL). Methanesulfonic acid (0.5 mL) was added and the reaction was stirred in the dark for twelve hours. The solvent was then removed *in vacuo* and the reaction mixture was purified by column chromatography.


The first fraction from the column was collected and, upon removal of the solvent, afforded a yellow oil. Acetone (10 mL) was added and the flask was swirled in order to dissolve the oil. After 20 min, white crystals of the *meso*-octabenzylcalix[4]pyrrole **6** (278 mg, 9.6% yield) had formed; they were collected and dried. ¹H NMR (CDCl₃) : 7.19 (m, 24H, ArH), 6.83 (m, 16H, ArH), 6.22 (br. s, 4H, NH), 5.62 (d, 8H, CH_(pyrrole)), 2.90 (m, 16H, ArCH₂). ¹³C NMR (CDCl₃) : 138.3, 134.1, 130.9, 127.7, 126.4, 106.5, 44.7. High Resolution negative FAB calc. for C₇₆H₆₇N₄ (1035.53674) found 1035.4720 (1.8 ppm).



Crystals of octabenzylcalix[4]pyrrole **6** suitable for X-ray study were grown from acetone. The conformation of the macrocycle in **6**·2(acetone)·acetone_{lattice} is 1,3-alternate (as in **1**·2(MeOH)), but the two acetone guests are both located on the *same* face of the ligand, held in place by single H-bonds (FIG 5). One of the acetone moieties lies much closer to the macrocycle than the other (N_{pyrrole}...O_{acetone} = 2.972(6) Å vs. 3.359(6) Å), reflecting an almost colinear (177(5)°) arrangement of N_{pyrrole}-H...O_{acetone} atoms in the former case. By contrast, the N-H...O angle formed with the more distant ketone measures 159(4)°. Octabenzylcalix[4]pyrrole may prove useful as a phase transfer catalyst for anions, or as a starting material for the synthesis of calix[4]pyrrole aryl-transition metal complexes which could be used as catalysts for various reactions or as a sensor for anions. This material is also a potential precursor to pyridine-containing systems (Example 6).


Tetraspirotetrahydrothiopyrancelix[4]pyrrole 7. Tetrahydrothiopyran-4-one (0.50g, 4.3 mmol) and pyrrole (0.288g, 4.3 mmol) were dissolved in ethanol (30 mL) and 10 drops of methanesulfonic acid were added. The mixture was stirred for 12 hours in the dark, after which time a white precipitate had formed. The solvent was removed *in vacuo* and the residue purified by column chromatography (SiO₂, CH₂Cl₂ eluant) affording the tetra sulfide macrocycle as a white powder (540 mg, 76 % yield). ¹H NMR (250 MHz, CD₂Cl₂) δ: 7.10 (s, 4H, NH), 5.97 (m, 8H, CH_{pyrrole}), 2.61 (m, 12H, CH_{pyran}) 2.28 (m, 12H, CH_{pyran}). ¹³C NMR (62.90 MHz,

CD₂Cl₂) δ : 136.0, 105.0, 39.5, 37.9, 24.9. High resolution FABMS calc for C₃₆H₄₄N₄S₄ 660.2449; found 660.2437 (Δ 1.7 ppm).




Crystals of macrocycle **7** suitable for analysis by X-ray diffraction techniques were grown by slow evaporation of a dichloromethane solution of the macrocycle (FIG. 6). Crystallographic data for **7**: Colorless crystals, triclinic, $P\bar{1}$, $Z = 2$, in a cell of dimensions: $a = 11.059(2)$, $b = 15.023(3)$, $c = 15.874(3)\text{\AA}$, $\alpha = 112.21(1)$, $\beta = 102.64(1)$, $\gamma = 103.07(2)^\circ$, $V = 2238.9(7)\text{\AA}^3$, $\rho_{\text{calc}} = 1.42\text{ g cm}^{-3}$, $F(000) = 998$. 3987 unique reflections, 2991 with $F_o > 4(\sigma(F_o))$. The final $R(F) = 0.0545$, $R_w(F^2) = 0.126$, goodness of fit = 1.133 for 336 parameters refined on F^2 . Calix[4]pyrrole **7** is a useful synthon for the synthesis of transition metal complexes on the periphery of the calixpyrrole which complexes are contemplated to be useful as sensors. Oxidation of macrocycle sulfide groups to sulfoxide or sulfone introduces further polarity into the macrocycle and renders it more water soluble, thus allowing use in biomedical applications (see Example 11). Additionally, the present inventors envision that analogous macrocycles containing NH, NH₂⁺, NHCH₃⁺, N(CH₃)₂⁺, NOH, NNH₂ and NN(CH₃)₂ functionalities instead of sulfur will be water soluble.

meso-Octanonylcalix[4]pyrrole 8. 10-Nonadecanone (3.14g, 11.12 mmol) and pyrrole (0.745g, 11.12 mmol) were dissolved in a mixture of ethanol (100 mL) and dichloromethane (50 mL). Methanesulfonic acid (20 drops) was added and the mixture was stirred for 12 hours. The solvent was then removed *in vacuo* and the product was purified by column chromatography on silica gel (hexane eluant) affording the product as a white powder (650 mg, 17 % yield). ¹H NMR (250 MHz, CDCl₃) δ : 6.99 (s, 4H, NH), 5.86 (s, 8H, CH), [1.77, 1.54, 1.21, 0.86 (br. multiplets, 152H, CH₂,CH₃)]. ¹³C NMR (62.90 MHz, CDCl₃) δ : 136.2, 104.6, 42.4, 37.7, 31.9, 30.3, 29.78, 29.75, 29.4, 24.0, 22.7, 14.1. High resolution FABMS calc for C₉₂H₁₆₅N₄ 1326.3034; found 1326.3015 (Δ 1.4 ppm).



Due to hydrophobic alkyl groups, calixpyrrole **8** is designed to have particular affinity for lipid bilayer membranes and is expected to serve as an insertion agent, i.e. insertion of macrocycle **8** is expected to allow passage of charged species (such as phosphorylated anti-viral agents) through cell membranes, thereby increasing the effectiveness of the agent (Example 11).

meso-Tetramethyl-mesotetraferrocenylcalix[4]pyrrole 9. Pyrrole (0.373 g, 5.56 mmol) and acetylferrocene (1.27 g, 5.56 mmol) were dissolved in methanol (50 mL) and stirred. Methanesulfonic acid (20 drops) was added and the reaction stirred for twelve hours. Triethylamine (30 drops) was then added to the reaction mixture followed by aspartic acid (2g) and the solvent removed *in vacuo*. The residue was purified by column chromatography (silica gel, CH_2Cl_2 eluant) yielding the product as a yellow foam (390 mg, 25% yield). ^1H NMR (CD_2Cl_2) 1.801-1.997 ppm (12H, m, CH_3), 3.931-4.198 ppm (36H, m, C_{10}H_9 (ferrocene)), 5.625-5.927 (8H, m), 7.929 (s, 4H, NH). ^{13}C NMR (CD_2Cl_2) 28.958, 31.016, 39.841, 39.857, 67.343, 67.525, 67.818, 69.031, 69.141, 69.356, 98.224, 104.588, 104, 932, 105.814, 136.866, 137.348, 137.482, 137.572



9

Calix[4]pyrrole **9** can be used as a reporter molecule or sensor for detecting anions or neutral molecules. The complexation of an anion in the calixpyrrole core will perturb the electronic states of the ferrocene moieties, thereby causing a shift in the ferrocene redox potential which can be detected by electrochemical techniques, including but not limited to cyclic voltammetry and Osteryoung-type square wave voltammetry.

While the present example demonstrates ester (**3**), acid (**4**), alkenyl (**5**), aryl (**6**), sulfide (**7**), and alkyl (**8**) substituents for calix[4]pyrroles, one of skill in the art would be able to

synthesize further meso-substituted calix[4]pyrrole macrocycles in light of the teachings herein and using standard organic chemical synthesis reactions as described herein.


Example 3

Novel β -Substituted Calix[4] pyrrole Macrocycles

Calix[4]pyrrole macrocycles with functional groups in the β -position (i.e., on the β -carbon of a pyrrole ring), and the syntheses thereof, are provided in the present example. Such β -substituted calixpyrroles may have substitution at one or more of the β -positions. To prepare such compounds of the present invention, either *meso*-octamethylcalix[4]pyrrole 1 is deprotonated and the substituent to be added at the β -position is then reacted with the activated calixpyrrole, or the calixpyrrole is synthesized from a β -substituted pyrrole.

Initially, sodium hydride was tried as a base to deprotonate the calixpyrrole. Sodium hydride was added to a solution of the macrocycle in DMF, after which bromoethylacetate was added in order to introduce an ester functionality to the calixpyrrole. However, this produced no reaction.

A stronger base (*n*-BuLi) was then tried; 4 equivalents of the base were added dropwise to a THF solution of the macrocycle at -78°C under an argon atmosphere. After 2 hours stirring at -78°C , 4 equivalents of bromoethylacetate were added and the solution was allowed to warm up to room temperature over 90 minutes. The THF was then removed *in vacuo* and the residue was purified using column chromatography on silica gel by eluting with CH_2Cl_2 . Several fractions were isolated from the column. The first proved to be unreacted calix[4]pyrrole. The second was the β -monosubstituted ester 10 and the third, the β -diester 11.




Calix[4]pyrrole β -mono- and disubstituted ester 10 and 11. *meso*-Octamethylcalix[4]pyrrole (1.0 g, 2.34 mmol; preparation described in Example 1) was dissolved with stirring in THF (100 mL, dry) under an argon atmosphere. The solution was

cooled to -78°C using a CO_2 /acetone bath. *n*-Butyllithium (5.83 mL, 9.34 mmol, 1.6 M in hexane) was added to the solution dropwise. After 2 hours, ethylbromoacetate (1.95 g, 11.68 mmol) was added dropwise to the solution. The reaction was then removed from the CO_2 bath and allowed to warm to room temperature. After 90 minutes, the solvent was removed *in vacuo* and EtOH (100 mL) was added to the flask. Water (100 mL) was slowly added with swirling, causing a white precipitate to form. The precipitate was collected by filtration, dried under high vacuum and then purified by column chromatography on silica gel eluting with CH_2Cl_2 . The monoester **10** (310 mg, 26%) and diester **11** (40 mg, 2.8%) were the second and third main products to be eluted from the column. **Mono-ester 10**: ^1H NMR (500 MHz, CD_2Cl_2) δ : 8.55 (s, 1H, NH), 7.15 (s, 1H, NH), 7.04 (s, 1H, NH), 7.00 (s, 1H, NH), 5.94 - 5.88 (overlapping multiplets, 5H, CH_{py}), 5.83 (m, 1H, CH_{py}), 5.72 (m, 1H, CH_{py}), 4.24 (q, $J = 7.15$ Hz, 2H, CH_2CH_3), 3.65 (s, 2H, CCH_2), 1.62-1.47 (overlapping singlets, 24H, CCH_3), 1.33 (t, 3H, $J = 7.15$ Hz). ^{13}C (DEPT) NMR (500 MHz, CD_2Cl_2) δ : 174.93 (C=O), 140.42 (C_{py}), 139.69 (C_{py}), 139.31 (C_{py}), 138.92 (C_{py}), 138.50 (C_{py}), 137.98 (C_{py}), 137.20 (C_{py}), 134.06 (C_{py}), 109.95 (CH_{py}), 107.07 (CCH_2COOEt), 103.41 (CH_{py}), 103.10 (CH_{py}), 102.90 (CH_{py}), 102.66 (CH_{py}), 101.37 (CH_{py}), 61.40 (CH_2CH_3), 37.16 (CCH_3), 35.50 (CCH_3), 35.43 (CCH_2), 35.17 (CCH_3), 33.95 (CCH_3), 29.20 (CCH_3), 29.05 (CCH_3), 28.90 (CCH_3), 28.68 (CCH_3), 14.47 (CH_2CH_3). High resolution FABMS calc for $\text{C}_{32}\text{H}_{42}\text{N}_4\text{O}_2$ 514.3308; found 514.3302 (Δ 1.2 ppm) **Bis-ester 11**: ^1H NMR (500 MHz, CD_2Cl_2) δ : 8.61 (s, 1H, NH), 8.52 (s, 1H, NH), 7.20 (s, 1H, NH), 7.00 (s, 1H, NH), 5.85 - 5.54 (several multiplets - overlapping, 4H, CH_{py}), 5.63 (d, 1H, CH_{py}), 5.55 (d, 1H, CH_{py}), 4.19 (m, 4H, CH_2CH_3), 3.60 and 3.59 (two singlets, total 4H, CCH_2), 1.58 (s, 6H, CCH_3), 1.52 (s, 6H, CCH_3), 1.49 (s, 6H, CCH_3), 1.44 (s, 6H, CCH_3), 1.29 (t, 2H, CH_2CH_3), 1.27 (t, 2H, CH_2CH_3). ^{13}C (DEPT) NMR (500 MHz, CD_2Cl_2) δ : 175.51 (C=O), 175.21 (C=O), 141.21 (C_{py}), 140.36 (C_{py}), 138.98 (C_{py}), 137.94, (C_{py}) 137.70, (C_{py}) 137.68 (C_{py}), 134.33 (C_{py}), 133.89 (C_{py}), 109.64 ($\text{C}_{\text{py}}\text{CH}_2$), 109.09 ($\text{C}_{\text{py}}\text{CH}_2$), 106.73 (CH_{py}), 105.08 (CH_{py}), 103.23 (CH_{py}), 102.90 (CH_{py}), 101.93 (CH_{py}), 100.72 (CH_{py}), 61.42 (CH_2CH_3), 37.21 (CCH_3), 36.89 (CCH_3), 35.52 (CCH_3), 35.19 (CCH_3), 33.97 (CCH_2), 33.93 (CCH_2), 29.05 (CCH_3), 28.90 (CCH_3), 14.41 (CH_2CH_3). High resolution FABMS calc for $\text{C}_{36}\text{H}_{48}\text{N}_4\text{O}_4$ 600.3676; found 600.3663 (Δ 2.1 ppm). This reaction may also be accomplished using other bases including but not limited to aryllithium or alkylolithium or in the presence of a coordinating ligand such as but not limited to tetramethylethylenediamine with an aryllithium or alkylolithium base. Alternately other electrophiles may be attached to the activated calixpyrrole macrocycle (e.g. CO_2 , ethylchloroformate, acrylonitrile). β -Substituted calix[4]pyrrole esters are useful as a reagent for the synthesis of other calixpyrrole derivatives as described herein.

Calix[4]pyrrole β -acid 12. Compound **10** was de-esterified to produce calix[4]pyrrole β -acid **12** by refluxing in a solution of NaOH in EtOH/ H_2O . The acid was clarified by adding

perchloric acid to the cooled reaction solution and then collected by filtration. This yielded a white powder.




Compound **10** (200 mg, 0.38 mmol) was stirred in 20 mL ethanol. The slurry was heated to reflux and NaOH_{aq} (20 mL, 2.0 M) was added. The reaction was heated at reflux until all the ester had been consumed (approximately 4 hours). The ethanol was then removed in vacuo and a further 50 mL of cold water was added to the solution. The solution was then acidified with concentrated perchloric acid to pH 1. The monoacid **12** then precipitated out as a white powder, was collected by filtration and dried under high vacuum to yield 146 mg (79%). ¹H NMR (500 MHz, CD₂Cl₂) δ: 7.84 (s, 1H, NH), 7.24 (s, 1H, NH), 7.01 (coincident singlets, 2H, NH), 5.90 - 5.76 (overlapping multiplets, 5H, CH_{py}), 3.71 (s, 2H, OCH₂), 1.51 (s, 6H, CCH₃), 1.50 (s, 6H, CCH₃), 1.48 (s, 6H, CCH₃), 1.47 (s, 6H, CCH₃). ¹³C NMR (500 MHz, CD₂Cl₂) δ: 179.13 (C=O), 139.96 (C_{py}), 139.20 (C_{py}), 139.09 (C_{py}), 138.60 (C_{py}), 138.52 (C_{py}), 137.45 (C_{py}), 133.96 (C_{py}), 109.27 (C_{py}), 107.11 (CH), 103.21 (CH), 103.14 (CH), 103.05 (CH), 102.91 (CH), 102.75 (CH), 102.14 (CH), 37.20 (CCH₃), 35.49 (CCH₃), 35.40 (OCH₂), 35.23 (CCH₃), 33.68 (CCH₃), 30.38 (CCH₃), 29.16 (CCH₃), 29.09 (CCH₃), 29.00 (CCH₃), 28.85 (CCH₃). High resolution FABMS calc for C₃₀H₃₈N₄O₂ 486.2989; found 486.2995 (Δ 1.1 ppm).

β-substituted “monohook” calix[4]pyrrole **12** is particularly useful for further substitution as described in Examples 8, 9, 10, and 11.

β-Octamethoxy-meso-tetraspirocyclohexylcalix[4]pyrrole 13. 3,4-Dimethoxypyrrole (0.568g, 4.5 mmol) and cyclohexanone (437 mg, 4.5 mmol) were dissolved in glacial acetic acid (10 mL), and the solution was stirred for 3 hours in the dark under argon, forming a black solution. The acid was then removed *in vacuo* and the residue purified by column chromatography (SiO₂, CH₂Cl₂ eluant). The product was collected as the second fraction from the column having an R_f = 0.33 (81 mg, 0.097 mmol) in a yield of 8.6%. Following chromatographic purification, the solvent was immediately reduced *in vacuo* and the product **13**

stored under argon in the freezer. ^1H NMR (300 MHz, CD_2Cl_2) δ : 6.11 (s, 4H, NH), 3.77 (s, 24H, OCH_3), 2.08 (s, br., 16H, CH₂), 1.45 (s, br., 24H). ^{13}C NMR (75.42 MHz, CD_2Cl_2) δ : 136.3, 75.73, 61.1, 42.5, 35.1, 26.3, 23.9. High resolution FABMS calc for $\text{C}_{48}\text{H}_{68}\text{N}_4\text{O}_8$ 828.5037 found 828.5025 (Δ 1.5 ppm).



5


β -Substituted calix[4]pyrrole **13** shows decreased anion binding affinity relative to compound **2** due to electron donating properties of the methoxy substituents. These types of compounds are useful as HPLC column media since low affinity constants and fast complexation-decomplexation rates generally lead to efficient separation of substrates (see Example 10).

10

β -Octabromo-meso-octamethylcalix[4]pyrrole 14. An oven-dried 100 mL round-bottomed flask containing a stir bar was charged with *meso*-octamethylcalix[4]pyrrole (0.69 g, 1.6 mmol) and 50 mL of dry tetrahydrofuran. With stirring, *N*-bromosuccinimide (2.34 g, 13.1 mmol) was added all at once, causing the reaction mixture to turn yellow. A reflux condenser and argon balloon were affixed to the flask, which was heated to gentle reflux in the dark. After 48 h, the clear red-brown solution was allowed to cool to room temperature, and the THF was removed *in vacuo*. The tan solid which remained was triturated with 20 mL of dichloromethane, and the CH_2Cl_2 solution was flash chromatographed on silica gel (1:1 CH_2Cl_2 -hexanes as eluent). Colorless-to-light-yellow fractions with $R_f = 0.73$ were combined and evaporated to afford a white solid. Drying under high vacuum gave 1.53 g (90%) of the desired product **14**. ^1H NMR (CDCl_3): δ 1.56 (s, 24H), 7.34 (br s, 4H). ^{13}C NMR (CDCl_3): δ 25.9, 38.2, 98.6, 130.4. LRMS (FAB⁺): m/z 1059, 979, 901, 821, 741. Elemental analysis calc. for **14**·2(DMSO) C, 31.61; H, 3.32; N, 4.61. Found C, 32.18; H, 3.31; N, 4.59.

15

20



Single crystals of compound **14** were grown by slow evaporation of a dilute dichloromethane solution of the macrocycle (FIG. 7). The structure reveals that compound **14** exists in a chair-like flattened 1,2-alternate conformation in the solid state (i.e., the dihedral angles between pyrrole rings and plane through the calixpyrrole *meso*-carbon atoms are 66.8°, 5.8°, -66.8° and -5.8°). Crystallographic summary for C₂₈H₂₈N₄Br₈: Large, colourless prisms were grown by slow evaporation from CH₂Cl₂. Crystals were monoclinic, P2₁/n Z = 2, in a cell of dimensions: a = 10.819(4), b = 12.042(4), c = 12.746(4) Å, β = 99.01(2)°, V = 1640.1(10) Å³, ρ_{calc} = 2.15 g cm⁻³, μ = 98.07 cm⁻¹, F(000) = 1008. A total of 7416 reflections were measured, 3763 unique (R_{int}(F²) = 0.091) on a Siemens P3 diffractometer using graphite monochromatised Mo Kα radiation (λ = 0.71073 Å) at -75 °C. The structure was refined on F² using SHELXTL/PC (ver 5.03, Siemens Analytical X-ray Instruments, Madison, WI, USA) to an R_w = 0.131, with a conventional R = 0.0575 (2231 reflections with F_o > 4[σ(F_o)]), and a goodness of fit = 1.105 for 182 refined parameters. The molecule lies on a crystallographic inversion centre at 0,0,0.

β-Substituted calix[4]pyrrole **14** shows increased anion binding affinity relative to the 'parent' calix[4]pyrrole **1** due to the electron withdrawing properties of the bromine substituents (see Example 10). Receptors with an increased anion binding ability such as **14** are useful as anion sequestering agents (e.g., in removal of phosphate and nitrate pollutants from aqueous environments) (see Example 12) or as synthons in reactions to produce further modified calix[4]pyrroles. The present inventors envision this system will be useful as a precursor to a calix[m]pyridino[n]pyrrole or a calix[m]pyridine.

While the present example describes ester (**10,11**), acid (**12**), alkoxy (**13**) and halo (**14**) β-substituents, one of skill in the art would be able to synthesize further modified β-substituted molecules in the light of the present disclosure and using standard organic chemical synthesis reactions as described herein.

β-Substituted calix[4]pyrrole macrocycles have similar utility to that of the *meso*-substituted molecules described herein.

Example 4

Use of Calix[n]arene in the Synthesis of Calix[n]pyrrole, and Pseudo Dimers Thereof

The present example provides the synthesis of a calix[n]arene-calix[n]pyrrole pseudo dimer. Reaction of *p*-*tert*-butylcalix[4]arene tetramethyl ketone (0.5 g, 0.57 mmol) with pyrrole (0.153 g, 2.29 mmol) in a dichloromethane/ethanol mixture, in the presence of methanesulfonic acid, followed by column chromatography on silica gel (CH₂Cl₂, eluant) afforded the calixpyrrole-calixarene pseudo dimer **15** in 32% yield. Compound **15** gave spectroscopic and analytical data in accord with the assigned structure. As a consequence of this template synthesis, the stereochemistry at the calixpyrrole *meso*-carbons is automatically defined. Thus this method of synthesis produces only one configurational isomer even though all four ketone groups involved in the synthesis of an individual calixpyrrole are asymmetrical (see Example 7).


Analytical and spectroscopic data for compound **15**: Analysis for C₇₂H₈₄N₄O₄: Calculated C 80.86%; H 7.92%; N 5.24%. Found C 80.57%; H 7.95%; N 5.11%. High resolution positive ion FABMS: Calculated (C₇₂H₈₄N₄O₄) 1068.649; Found 1068.648 (Δ 1.4 ppm). ¹H NMR (CD₂Cl₂) δ : 11.22 (s, 4H, NH), 7.22 (s, 8H, ArH), 5.83 (d, J = 2.5 Hz, 8H, CH), 4.35 (d, J = 12.5 Hz, 4H, ArCH₂Ar), 3.80 (s, 8H, OCH₂), 3.37 (d, J = 12.5 Hz, ArCH₂Ar), 1.58 (PyCCH₃), 1.18 (s, 36H, *tert*-Bu). ¹³C NMR (CDCl₃) δ : 151.3, 146.8, 137.2, 134.4, 126.1, 101.7, 90.2, 37.9, 34.1, 31.4, 29.6, 21.1.

Crystals were obtained by slow evaporation of a dichloromethane solution of compound **15**. The crystals were then subject to X-ray diffraction structural analysis. The resulting structure shows the calixarene in a cone conformation. The pyrrole NH groups are within hydrogen bonding distance of the lower rim calixarene oxygen atoms (FIG. 8).

Crystallographic summary for **15**.xCH₂Cl₂. Large regions of disordered solvent of unknown stoichiometry in this structure presented difficulties with this analysis, however the structure of the calixarene-calixpyrrole pseudo dimer is a reliable description of this unique and interesting molecule. Faintly yellow crystals, monoclinic, *C*2/*m*, *Z* = 4 in a cell of dimensions *a* = 25.836(5) Å, *b* = 20.139(3) Å, *c* = 15.46(2) Å, β = 106.0(1)°, *V* = 7719(2) Å³. ρ_{calc} : *x* = 4, 1.21 gcm⁻³; *x* = 6, 1.36 gcm⁻³. 9112 unique reflections of which 4121 were observed (*I* > 2 σ (*I*)). Data collected at -90°C on a Siemens P3 diffractometer. The final *R* = 0.148.

The ¹H NMR spectrum of **15** in dichloromethane-*d*₂ shows that the pyrrole NH protons, which are normally observed at around 7 ppm in other calix[4]pyrroles, resonate at 11.22 ppm (FIG. 9). This can be attributed to hydrogen bonding between the pyrrole NH groups and the calixarene oxygen atoms. CD₃OD was added to the NMR solution in an attempt to disrupt this hydrogen bonding interaction, however this had a negligible effect on the ¹H NMR spectrum. Similarly, addition of 10.0 equiv. tetrabutylammonium fluoride (which has been shown to bind strongly to other calix[4]pyrroles) caused no changes in the NMR

spectrum. Based on these findings, the present inventors conclude that the dimer is adopting a cylindrical conformation in solution due to these clearly favorable NH - O hydrogen bonds.




5

These types of calixarene-calixpyrrole molecular cylinders may be useful as receptors of neutral guests, as they possess a hydrophobic cavity. Additional functional groups could be appended to the free phenolic OH groups on the calixarene. These may be used to attach the molecule to a polymer or other solid support. Such solid supported compounds may be used in column chromatography, for example. Alternatively, an optical or electrochemical sensing group such as anthracene or ferrocene could be attached and may be able to sense the inclusion of neutral species in the cylinder and therefore act as a sensor. These materials may also be able to extract toxic organic substances (such as insecticides and PCBs) from aqueous solutions and, being potential receptors for zwitterions, aid in the purification of amino acids, peptides, proteins, and the like.

15

In addition to a dimer cylinder, trimers and larger tubular polymers may be prepared. The synthesis of a calixarene/calixpyrrole/calixarene pseudo trimer 16 is illustrated below:



Calix[4]arene tetra ketones in the 1,3-alternate conformation may be used to make calixarene-calixpyrrole tubular polymers which may have interesting properties, either alone or with a guest species bound within the polymer tube.

5

Example 5

Synthesis of Calix[n]pyrroles

Synthesis of Calix[n]pyrroles where n = 4, 5, 6, 7, or 8. Calix[n]pyrroles where n is greater than 4 have not previously been described. The present example provides methods for synthesis of higher order cyclic oligomers where n is 4 - 8. Prior art methodology for preparing calix[4]pyrroles involved the condensation of a ketone with a pyrrole using a Brønsted acid such as methanesulfonic acid or hydrochloric acid. This method produces exclusively calix[4]pyrrole. The present inventors have developed a methodology that employs Lewis acid catalysis of the condensation of pyrroles and ketones to yield calix[4]pyrrole and the previously undisclosed calix[5, 6, 7, or 8]pyrroles.

15

Lewis acids have previously been used to catalyze porphyrinogen formation. The present inventors have extended this methodology to calix[4]pyrrole formation. A general synthetic protocol for the preparation of calix[n]pyrroles at ambient temperature employing a variety of Lewis acid catalysts is also described herein and allows a high yield and purity of the calix[n]pyrrole product. A pyrrole and a ketone may undergo Lewis acid-catalyzed condensation to form a generalized calix[n]pyrrole having structure I. The reaction conditions have been optimized for the preparation of higher calix[n]pyrroles (n>4) by varying pyrrole and

20

the ketone ratios and employing $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as a catalyst. Optimal conditions for the formation of higher order calix[n]pyrroles ($n > 4$) are a 1 : 1 to a 1.25 : 1 molar ratio (ketone:pyrrole) with a 10% Lewis acid catalyst. The concentration of starting compounds ranged from 50 mM to 1 mM depending on the solvent used e.g. dichloromethane (dry), chloroform and acetonitrile, although it may be possible to use other solvents or solvent mixtures to obtain optimum yields of a given calix[n]pyrrole.

Lewis acids such as $\text{BF}_3 \cdot \text{OEt}_2$, BF_3 -methanol, VOCl_3 , TiCl_4 , and $\text{Mn}(\text{OAc})_3$ have been used as catalysts and it has been found that TiCl_4 in dry dichloromethane produces higher yields of calix[n]pyrroles than the other catalysts. Lewis acids may also be used in doped zeolites, monomorphonites, clays, silica and alumina which can provide a convenient site for calix[n]pyrrole formation.

The conditions for the preparation of calix[n]pyrroles ranging from $n = 4$ to 10 are: acetone (0.5808 g, 10 mmol) and pyrrole (0.67 g, 10 mmol) were dissolved in dry dichloromethane (150 mL) under an argon atmosphere with stirring under argon for 15 minutes. The catalyst, $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.5 mL of 1M solution in dichloromethane, 0.5 mmol), was slowly added *via* syringe. The reaction mixture was stirred under argon for between 24 to 48 hours. HCl (3 mL of 3 %) was then added and the mixture stirred for 10 minutes. The reaction mixture was transferred to a separatory funnel and the organic phase was washed with water and dried over sodium sulfate. The solvent was evaporated and the crude reaction mixture was filtered over a short neutral alumina column using dichloromethane as the mobile phase. The calix[n]pyrrole products were purified by flash chromatography or by preparative HPLC using dichloromethane-hexane (3:1) a mobile phase. Yield 78 %. The ratio for n is as follows: 1, $n = 4$, 60 %; 17, $n = 5$, 12 %; 18, $n = 6$, 9%; 19, $n = 7$, 5%; 20, $n = 8$, 3%. Separated calix[n]pyrroles were characterized by HRMS: Cyclic pentamer: 17, $n = 5$, For $\text{C}_{35}\text{H}_{45}\text{N}_5$ calc. 535.367497, found. 535.367194. For $[\text{MH}^+]$ $\text{C}_{35}\text{H}_{46}\text{N}_5$ calc. 536.375322, found 536.374278. Cyclic hexamer: 18 $n = 6$ For $\text{C}_{42}\text{H}_{54}\text{N}_6$ calc. 642.440996, found 642.440390. Cyclic heptamer: 19 $n = 7$ For $\text{C}_{49}\text{H}_{63}\text{N}_7$ calc 749.514496, found 746.514133. Cyclic octamer: 20 $n = 8$ For $\text{C}_{56}\text{H}_{72}\text{N}_8$ calc. 856.58796, found 856.58

Chloroform and acetonitrile may also be used as solvents. $\text{BF}_3 \cdot \text{MeOH}$ was used as a catalyst with 73 % overall yield. When 1-alkyl, or 1-aryl pyrrole were used, the reaction again produced cyclic systems; *N*-substituted calix[n]pyrroles where n is from 4 to 10, with the main product $n = 4$, but with higher ratio for $n > 4$ than in case of pyrrole (the ratio estimate is 35: 12: 9: 7 : 5) for $n = 4 - 8$, for 1-methylpyrrole and ratio 29:12:9:8:6 for 1-phenylpyrrole.


The procedure for the use of high valent Lewis acid catalysts (e.g. TiCl_4) in calix[n]pyrrole synthesis is as follows: acetone (0.581 g , 10 mmol) and pyrrole (0.67 g, 10 mmol) were dissolved in 100 mL of dry dichloromethane under argon. Titanium tetrachloride (1 mL of 1M solution in dichloromethane, 1 mmol) was then added to the solution under argon while stirring at room temperature. Immediately after addition, a precipitate was formed, which


redissolved after 5 minutes stirring. The reaction mixture was stirred at room temperature for 20 hours, then washed with 3 % HCl, water and dried over sodium sulfate. The product was purified by column chromatography on neutral alumina. Final separation was performed by flash chromatography on silica gel using dichloromethane-hexane 4:1. The overall yield was 67% with a calix[n]pyrrole distribution as follows: 88 (1, n = 4), 8 (17, n=5), 6 (18, n=6), 3 (19, n = 7), 2 (20, n=8).

The procedure has been repeated using analogous conditions to those used with titanium tetrachloride but using other Lewis acids such as VOCl_3 , $\text{Mn}(\text{OAc})_3$, SnCl_4 , AlCl_3 , or $\text{UO}_2(\text{OAc})_2$ and also using a number of different solvents.


10 A variety of cation exchange resins (Dowex) were employed in their H^+ rich forms, however, the product was exclusively the cyclic tetramer.

Synthesis of Calix[n]pyrroles using clays. Montmorillonite K10 (Aldrich) was added to a 250 mL flask and dried at 120°C / 0.2 Torr for 2 hours. After cooling to room temperature under argon, dry dichloromethane (100 mL) was added, followed by acetone (0.0581 g, 1 mmol) and pyrrole (0.067g, 1 mmol) successively. The reaction mixture (suspension) was vigorously stirred under argon for 8 hours. The solid material was filtered through Celite and washed with dichloromethane (25 mL). The combined dichloromethane filtrates were evaporated and the product was purified on a short neutral alumina column (dichloromethane eluant). The products separated by preparative HPLC on a silica gel column using dichloromethane-hexane as the mobile phase. Overall yield was 68% with the main product being calix[4]pyrrole 1 (59%). Additionally, calix[5]pyrrole 17 (6% yield and 3% of higher oligomers ($n > 6$)) were isolated.







17



18




19



20

Synthesis of calix[n]pyrroles via cyclization of 2-[(dialkylhydroxy)methyl]pyrrole

Calix[n]pyrroles may also be synthesized from 2-[(dialkylhydroxy)methyl]pyrroles. These molecules are cyclized using either a Brønsted acid or a Lewis acid. One approach for the preparation of 2-[(methylalkyl)hydroxymethyl]pyrrole derivative is based on the reaction of commercially available 2-acetylpyrrole with alkylmagnesiumbromides (Grignard reagents) affording the desired product **21** (R = alkyl) as a starting building block for Brønsted acid, or Lewis acid catalyzed cyclization.





10

2-Acetylpyrrole (0.109 g, 1 mmol) was dissolved in 50 mL of dry THF under argon. The solution was then cooled to -78°C . A solution of MeMgBr (1 mL of a 1 M solution in

diethylether) was added via syringe while stirring. The reaction mixture was allowed to warm up to -15°C for 10 minutes. It was then cooled down and another 1.5 molar equivalents MeMgBr (1.5 mL of 1M solution; 1.5 mmol) was slowly added at -78°C . The reaction mixture was stirred at that temperature for 15 minutes and then slowly allowed to warm up to 0°C .
5 After stirring for two hours at 0°C , the reaction was quenched with MeOH and evaporated to dryness. The residue was redissolved in dichloromethane (50 mL) and washed with a 5% solution of ammonium chloride and water. The 2-(dimethylhydroxymethyl)pyrrole was subjected to acid and Lewis acid catalytic macrocyclization giving mixture of calix[n]pyrroles at 65% yield with $n = 4-8$ as the products with the main product being 1. The present inventors
10 envision this synthesis could be extended to pyrroles with ketone groups in the α -position.

β -Octaethylcalix[4]pyrrole 22. 3,4-Diethylpyrrole (0.369 g, 3 mmol) was dissolved in dry THF (30 mL) under argon and the solution was cooled to -78°C . A MeMgBr solution (3 mL of 1 M solution in diethylether, 3 mmol) was slowly added and stirred at that temperature for 10 minutes. The solution was allowed to warm up to -10°C for 10 minutes and was then
15 cooled to -78°C . At this point, 1.2 molar equivalents of *tert*-BuLi was added (3.6 mL of a 1 M molar solution in hexane, 3.6 mmol) and the reaction mixture was kept at -78°C for 30 minutes. The reaction mixture was allowed to warm up to -10°C for 10 minutes and then cooled to -78°C . Acetone (0.29 g, 5 mmol) was added to the solution at that temperature and stirred for 30 min. at -78°C , then allowed slowly warm up to -10°C and kept at that temperature for 1 hour. The
20 reaction was quenched by addition of 1mL of saturated water solution of ammonium chloride. After evaporation of the organic solvent, the product 21 ($R_1 = \text{Me}$, $R_2 = R_3 = \text{Et}$) was extracted into dichloromethane and isolated by flash chromatography on silica gel with dichloromethane as the mobile phase. Yield 78%. HR MS for $\text{C}_{11}\text{H}_{19}\text{NO}$ calc. 181.146664, found 181.14781. 2-[(Dimethyl)hydroxymethyl]-3,4-diethylpyrrole 21 was subjected to acid
25 and Lewis acid catalysed cyclization. The mixture of calix[n]pyrroles was isolated in 45% net yield. The main product has $n = 4$ 22 : $(\text{C}_{11}\text{H}_{19}\text{N})_4$ found 652; but a 5% total yield of higher analogues was also obtained.





Pyrrole can be substituted for 3,4-diethylpyrrole in this procedure.

The present inventors envision that 1-alkyl pyrrole and 1-alkyl-3,4-diethylpyrrole and ketones could be used in a modification of the above procedure: 1-methylpyrrole or 1-methyl-3,4-diethylpyrrole (3 mmol) are dissolved in dry THF (30 mL) under argon and cooled to -78°C. 1.2 molar equivalents of *tert*-BuLi are added (3.6 mL of a 1 M molar solution in hexane, 3.6 mmol) and the reaction mixture kept at -78°C for 30 minutes. The reaction is warmed to -10°C for 10 min. and then cooled to -78°C. A general ketone (5 mmol) added and the reaction mixture was stirred for 30 min. at -78°C. The solution is allowed to slowly warm up to -10°C and stirred at that temperature for 1 hour. The reaction is quenched with 1 mL of a saturated water solution of ammonium chloride. After evaporative removal of the organic solvent, the product is extracted to dichloromethane and isolated by flash chromatography on silica gel using dichloromethane as a eluent. The products: 2-[(dialkylhydroxy)methyl]pyrrole derivatives were used for the preparation of calix[n]pyrroles using the methodology described above.

Example 6

Conversion of Calix[m + n]pyrroles to Calix[m]pyridino[n]pyrrole

A general synthetic method for the preparation of calix[m]pyridino[n]pyrroles where m and n designate a number of pyridines and pyrroles in the macrocycle, respectively, is unknown to date. Calix[2]pyridino[2]pyrroles and calix[1]pyridino[3]pyrroles have been previously synthesized *via* an organometallic mediated transformation using zirconium and other metals. However, calix[m]pyridino[n]pyrroles where m > 2 have not been reported.

The synthetic approach provided by the present inventors differs from the prior art methods and provides access to a broad variety of calix[m]pyridino[n]pyrroles and calix[m]pyridines. The approach of the present synthetic protocol is based on a non organometallic-mediated pyrrole ring expansion involving one or more of the pyrrole rings contained in a calix[m+n]pyrrole macrocycle.

Expansion of pyrrole within an existing calix[m+n]pyrrole involves the cycloaddition of dichlorocarbene to a particular pyrrolic subunit followed by rearrangement to yield a corresponding pyridine. Dichlorocarbene may be generated under basic or neutral conditions.

One or more pyrrolic subunits may react in this procedure specifically, varying the ratio of dichlorocarbene to calix[m+n]pyrrole allows control of the extent of conversion of pyrrole to pyridine which, in turn, determines the ratio of pyridine subunits to pyrrole subunits in the calix[m]pyridino[n]pyrroles.

The present inventors have also discovered a solvent dependence for the conversion of calix[m+n]pyrroles to calix[m]pyridino[n]pyrroles. The use of dioxane leads mainly to one and two converted units, while the use of 1,2-dimethoxyethane and prolonged reaction time provides for the full conversion of calix[m+n]pyrroles to calix[m]pyridines. Partial conversion of calix[m+n]pyrroles to calix[m]pyridino[n]pyrroles is also accomplished by the present methods.

Additionally, the present inventors have found that dichlorocarbene generated under neutral conditions (from sodium salts of trichloroacetic acid, for example) gives better yields than generating the dichlorocarbene under basic conditions (i. e. from the reaction of chloroform with sodium ethoxide).






A first method utilizes dichlorocarbene generated under neutral conditions and achieves fully converted calixpyridine product. *meso*-Octamethylcalix[4]pyrrole (10 mmol) was dissolved in dry 1,2-dimethoxyethane (75 mL). Four molar equivalents of dry sodium trichloroacetate were added per pyrrole ring. The reaction mixture was heated to reflux under an argon atmosphere for 20 hours. After cooling to room temperature, the solution was filtered and the organic solvent was evaporated *in vacuo*. The resulting residue was redissolved in dichloromethane, treated with charcoal, filtered over celite and evaporated to dryness. The crude product was purified over a short neutral alumina column (dichloromethane eluant). Final separation was achieved by flash chromatography on silica gel with dichloromethane and methanol (0-10 %) as the mobile phase. Yield of fully converted product **27** is 35-50 %. HRMS: For $C_{32}H_{33}N_4Cl_4$ $[MH^+]$: calc. 613.145933, found 613.145490.

A second method employs an incomplete calix[4]pyrrole expansion process to form mixed calix[m]pyridino[n]pyrroles. Mixed systems were obtained using 1-3 molar equivalents of sodium trichloroacetate for each pyrrole ring in the macrocyclic system. The molar ratio of pyrrole/dichlorocarbene allows one to "tune" the number of pyrroles that are converted to pyridines. Using one molar equivalent of sodium trichloroacetate for each pyrrole unit, the main product was calix[1]pyridino[3]pyrrole **23** (76% yield). Using 2 equivalents, the main product was a mixture of the two calix[2]pyridino[2]pyrrole isomers **24** and **25** (83% yield). Using 3 molar equivalents, calix[3]pyridino[1]pyrrole **26** was obtained (56% yield). These compounds were isolated by flash chromatography and separation was obtained on semipreparative HPLC with dichloromethane-acetonitrile as a solvent and a gradient of

acetonitrile from 0-5%. After column and HPLC separation, the following 3-chloropyridine macrocyclic products were characterized by HRMS: Monoconversion: HRMS For $C_{29}H_{35}N_4Cl$ calc. 474.255025, found 474.254879. Bis conversion: HRMS For $C_{30}H_{34}N_4Cl_2$ calc. 520.216053; found 520.215789 and 512.215985. For tris conversion: HRMS For $C_{31}H_{33}N_4Cl_3$ calc. 566.177081, found 566.178902.

Mixed molecule systems were further obtained using a third procedure. Dioxane was used as a solvent under the conditions described which resulted in full conversion of the calix[4]pyrrole to calix[4]pyridine, the main products were calix[1]pyridino[3]pyrrole **23**, calix[2]pyridino[2]pyrrole **24** and **25** and calix[3]pyridino[1]pyrrole **26**. Conditions: reflux under argon for 15 hours, purification and separation of products as above.


Sodium (1.15 g, 50 mmol) was dissolved in absolute ethanol (25 mL) in a flask fitted with a reflux condenser and a mechanical stirrer and the flask was flushed with argon. Calix[4]pyrrole (15 mmol) in chloroform (5 mL) was added and mixture was heated to 55°C for 18 hours. The sodium chloride was filtered off, the residue was treated with charcoal and purified on a short neutral alumina column followed by flash chromatography on silica gel with dichloromethane-methanol (from 0-5%). The main product, was isolated as unreacted starting calix[4]pyrrole with only 3-9% conversion.

**23****24****25****26****27**

Example 7





Configurational Isomers of Calixpyrrole

The structures of calix[4]pyrroles described herein are drawn two dimensionally. This representation obscures the fact that calixpyrroles containing more than one asymmetrically-substituted *meso* position will be formed as a mixture of configurational isomers. For example, compound 5 (see Example 2), contains four asymmetrically substituted *meso* positions and can exist as four different isomers which are chemically distinct from each other and may be separated *via* HPLC chromatographic techniques. A two dimensional representation is as follows.



10

The four possible configurational isomers of compound 5 are as follows.



Therefore, as used herein, all two dimensional representations of calixpyrroles containing more than one asymmetrically substituted *meso*-position refer to all possible isomers ($\alpha, \alpha, \alpha, \alpha$), ($\alpha, \alpha, \alpha, \beta$), ($\alpha, \beta, \alpha, \beta$) and ($\alpha, \alpha, \beta, \beta$) unless otherwise stated. For calix[5]pyrrole there will also be four possible configurational isomers ($\alpha, \alpha, \alpha, \alpha, \alpha$), ($\alpha, \alpha, \alpha, \alpha, \beta$), ($\alpha, \beta, \alpha, \beta, \alpha$) and

15

($\alpha,\alpha,\alpha,\beta,\beta$), however for $n>5$ the number of possible configurational isomers will increase above four.

5


Example 8

Calix[n]pyrrole Derivatives, Multimers, Conjugates, and Synthesis Thereof


Calix[n]pyrrole derivatives, multimers and conjugates have been synthesized. Such molecules possess useful binding properties for anion, chiral substrate, or neutral molecule separation, for salt binding, for acting as sensors for anions or neutral guests and the like.

10 Amido-calixpyrroles **28** and **29** are useful as anion binding agents. Additionally they serve as model compounds for calixpyrrole modified silica gels wherein the calixpyrrole is linked to the modified silica *via* amide bonds.


Mono- β -butylamidocalix[4]pyrrole 28. Compound **12** (' β -hook' acid) (300 mg, 0.62 mmol) was dissolved in DMF (20 mL, dry) under and argon atmosphere. *n*-Butylamine (45 mg, 0.62 mmol) was added followed by BOP PF₆ (327 mg, 0.74 mmol) and triethylamine (125 mg, 1.2 mmol). The reaction mixture was stirred for 48 hours. The DMF was then removed in vacuo and the product purified by column chromatography (SiO₂, CH₂Cl₂: CH₃OH 99:1 eluant) affording compound **28** as a white foam (331 mg, 99%). ¹H NMR (250 MHz, CD₂Cl₂) δ : 8.80 (s, 1H, NH), 7.24 (s, 1H, NH), 7.01 (coincident s, 2H, NH), 5.92-5.82 (m, 7H, 6CH_{py} and NH_{amide}), 5.70-5.69 (1H, d, CH_{py}), 3.51 (s, 2H, OCH₂), 3.29 (q, J = 6.2 Hz, 2H, CH₂), 1.66-1.30 (coincident resonances, 28H, 8CH₃ *meso*, 2CH₂ *hook*), 0.96 (t, 3H, CH₃ *hook*). ¹³C NMR (62.9 MHz, CD₂Cl₂) δ : 173.0, 140.2, 139.7, 139.2, 138.9, 138.4, 138.1, 137.4, 134.2, 110.7, 106.9, 103.2, 103.2, 103.1, 102.7, 102.5, 101.6, 39.7, 37.1, 35.9, 35.5, 35.4, 35.2, 32.1, 29.2, 29.0, 28.7, 20.5, 14.0. High resolution FABMS calc for C₃₄H₄₇N₅O 541.3781; found 25 541.3782 (Δ -0.2 ppm).



Mono-meso-butylamidocalix[4]pyrrole 29. Compound **4** ('*meso*-hook' acid) (268 mg, 0.43 mmol) was dissolved in DMF (20 mL, dry) under and argon atmosphere. *n*-Butylamine (31.6 mg, 0.43 mmol) was added followed by BOP PF₆ (230 mg, 0.52 mmol) and triethylamine (87 mg, 0.86 mmol). The reaction mixture was stirred for 48 hours. The DMF was then removed in vacuo and the product purified by column chromatography (SiO₂, CH₂Cl₂: CH₃OH 99:1 eluant) affording compound **29** as a white foam (118 mg, 41%). ¹H NMR (250 MHz, CD₂Cl₂) δ : 7.26 (s, 2H, NH), 7.17 (s, 2H, NH), 5.93 (s, 2H, CH_{py}), 5.92 (s, 2H, CH_{py}), 5.90




Mono-β-octadecylamidocalix[4]pyrrole 30. 'β-hook' acid **12** (250 mg, 0.51 mmol) was dissolved in DMF (20 mL, dry) under and argon atmosphere. Octadecylamine (139 mg, 0.51 mmol) was added followed by BOP PF₆ (273 mg, 0.61 mmol) and triethylamine (105 mg, 1.03 mmol). The reaction mixture was stirred for 72 hours. The DMF was then removed *in vacuo* and the product purified by column chromatography (SiO₂, CH₂Cl₂: CH₃OH 99:1 eluant) affording the product **30** as a yellow oil (330 mg, 87%). ¹H NMR (250 MHz, CD₂Cl₂) δ: 8.79 (s, 1H, NH_{py}), 7.50 (s, 1H, NH_{py}), 7.03 (s, 2H, NH_{py}), 5.89-5.80 (coincident multiplets, 7H, 6CH_{py} and NH_{amide}), 5.67 (d, 1H, NH_{py}), 3.49 (s, 2H, CH₂CO), 3.26 (q, J = 6.4 Hz, 2H, CH₂), 1.68 (m, 12H, 4CH₃), 1.49 (m, 12H, 4CH₃), 1.29 (m, 32H, CH₂), 0.90 (t, J = 6.8 Hz, 3H, CH₃). ¹³C NMR (62.90 MHz, CD₂Cl₂) δ: 173.0, 140.2, 139.7, 139.2, 139.0, 138.4, 138.1, 137.4, 134.3, 110.8, 107.0, 103.3, 103.2, 102.7, 102.5, 101.7, 40.1, 37.2, 35.9, 35.50, 35.45, 35.2, 32.4, 30.2, 30.1, 29.8, 29.2, 29.0, 28.7, 27.4, 23.1, 14.3. High resolution FABMS calc for C₄₈H₇₅N₅O 737.5972; found 737.5957 (Δ 2.0 ppm). The amide NH group of this receptor contributes to the calix[4]pyrrole anion binding process by providing a fifth hydrogen bond to the bound anion (see Example 10)




25 Molecules such as **30** are designed for insertion into lipid bilayers and are expected to be useful for transport.

Propyl bridged β -amidocalix[4]pyrrole dimer 31. A calixpyrrole amide dimer compound **31** was formed by coupling 1,3-diaminopropane to the calixpyrrole acid **12**. Again, this amide was formed as a DMF complex. This type of receptor is expected to be an effective host for poly-anionic guest species.




Compound **12** (100 mg, 0.206 mmol) and 1,3-diaminopropane (7.62 mg, 0.103 mmol) were dissolved in 5 mL DMF under an argon atmosphere. BOP (109 mg, 0.247 mmol) and Et_3N (41 mg, 0.411 mmol) were added in 10 mL dry DMF, and the solution was stirred overnight. The solvent was then removed under high vacuum and the residue was purified by column chromatography (SiO_2 , CH_2Cl_2 - 1.0% MeOH), affording the dimer **31** as a DMF complex (65 mg, 62%). ^1H NMR (500 MHz, CD_2Cl_2) δ : 8.63 (s, 2H, NH), 7.93 (DMF), 7.35 (s, 2H, NH), 7.01 (s, 2H, NH), 6.35 (t, 2H, NH_{amide}), 5.88 - 5.67 (overlapping multiplets, 14H, CH_{py}), 3.51 (s, 4H, CCH_2), 3.25 (m, 4H, CH_2), 2.90 (DMF), 2.82 (DMF). High resolution FABMS calc for $\text{C}_{63}\text{H}_{82}\text{N}_{10}\text{O}_2$ 1010.6622; found 1010.6630 (Δ -0.7 ppm). This methodology can be extended to produce calixpyrrole trimers, networks of calixpyrroles, linear strings of calixpyrroles, or dendrimer type arrays of calixpyrroles. Additionally all these multimers could be synthesized using the *meso*-“monohook” acid **4** in a similar synthetic scheme as described herein for the β -substituted molecules.

β -Amidobenzo-15-crown-5-meso-octamethylcalix[4]pyrrole 32. 4'-Aminobenzo-15-crown-5 has been coupled to calix[4]pyrrole β -monoacid **12** to produce a ditopic receptor which is capable of binding a whole salt (i.e. $\text{Na}^+ + \text{Cl}^-$) (FIG. 10) or a zwitterionic amino acid. The calixpyrrole provides an anion binding site and the crown ether a cation binding site, thereby providing a salt complexing agent.




meso-Octamethylcalix[4]pyrrole β -monoacid **4** (281 mg, 0.58 mmol) and 4'-aminobenzo-15-crown-5 (164 mg, 0.58 mmol) were dissolved in dry DMF (20 mL) and stirred under an argon atmosphere. Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP PF₆) (0.306 g, 0.69 mmol) and triethylamine (116 mg) in 20 mL DMF were added and the reaction mixture stirred overnight. The solvent was then removed under high vacuum and the residue purified by column chromatography (SiO₂, CH₂Cl₂ - 1 to 2% MeOH) affording the amide **32** as a white foam containing alkali metal cations (0.463g). ¹H NMR (250 MHz, CD₂Cl₂) δ : 8.74 (s, 1H, NH_{py}), 7.76 (s, 1H, NH_{am}), 7.45 (s, 1H, ArH), 7.34 (s, 1H, NH_{py}), 7.09 (s, 1H, NH_{py}), 7.02 (s, 1H, NH_{py}), 6.87 (m, 2H, ArH), 4.10 (m, 4H, 2OCH₂), 3.87 (m, 4H, 2OCH₂), 3.71 (m, 10H, 4OCH₂ + CCH₂CO), 1.59 (s, 6H, 2CH₃), 1.57 (s, 6H, 2CH₃), 1.51 (s, 12H, 4CH₃). ¹³C NMR (62.5 MHz, CD₂Cl₂) δ : 171.48, 149.52, 145.96, 140.08, 139.56, 139.18, 138.78, 138.31, 138.14, 137.67, 134.48, 132.73, 114.66, 112.44, 109.97, 107.19, 106.91, 103.26, 103.13, 103.07, 102.83, 102.31, 101.71, 70.94, 70.40, 69.73, 69.57, 69.04, 37.12, 36.82, 35.45, 35.41, 35.19, 29.17, 28.97, 28.79. FABMS calc for C₄₄H₅₇N₅O₆ 751.4309; found 751.4310 (Δ -0.2 ppm). Further cation binding agents that may be coupled to calix[n]pyrrole include polyethers; texaphyrins and cryptands for example. Anion binding agents such as sapphyrin or other expanded porphyrins such as rubyrin, rosarin, amethyrin and turcasarin may also be appended to produce polyanionic receptors. Additionally neutral binding groups such as hydrogen bonding arrays may be appended to the calix[n]pyrrole.

Mono-meso-ferroceneamidocalix[4]pyrrole 33. The synthesis of new molecular devices designed to sense and report the presence of a particular substrate is an area of analytical chemistry which is attracting intense current interest. Ferrocene is a stable organometallic moiety that has been incorporated into *meso*- and β -pendant arms of calix[4]pyrrole macrocycles forming molecules which may be used as electrochemical sensors.



- Compound 4 ('meso-hook' acid) (325 mg, 0.52 mmol) (see Example 2) was dissolved in DMF (20 mL, dry) under an argon atmosphere. Methylaminoferrocene (112 mg, 0.52 mmol) was added followed by BOP PF₆ (278 mg, 0.63 mmol) and triethylamine (106 mg, 1.2 mmol). The reaction mixture was stirred for 48 hours. The DMF was then removed *in vacuo* and the product purified by column chromatography (SiO₂, CH₂Cl₂: CH₃OH 99:1 eluant) affording an orange foam **33** (288 mg, 67%). ¹H NMR (250 MHz, CD₂Cl₂) δ: 7.24 (s, 2H, NH), 7.16 (s, 2H, NH), 5.93 (d, 4H, CH_{py}), 5.89 (d, 4H, CH_{py}), 5.61 (br.t, 1H, NH_{amide}), 4.16 (s, 7H, FcH + FcCH₂), 4.13 (m, 2H, FcH), 4.07 (m, 2H, FcH), 2.05 (t, 2H, CH₂), 1.94 (coincident resonances, 16H, 2CH₂ hook + 6CH₂ cyclohexyl, 1.45 (br., 23H, CH₂ hook + CH₃ + 9CH₂ cyclohexyl). ¹³C NMR (62.90 MHz, CD₂Cl₂) δ: 172.0, 137.6, 137.1, 137.0, 136.8, 104.0, 103.9, 103.8, 103.6, 85.7, 68.9, 68.5, 68.4, 40.0, 39.8, 39.5, 38.9, 38.8, 37.2, 37.1, 37.0, 36.9, 36.7, 26.4, 25.6, 23.2, 23.1, 20.9. High resolution FABMS calc for C₅₁H₆₃N₅O⁵⁶Fe 817.4382; found 817.4364 (Δ 2.2 ppm).
- Mono-β-ferroceneamidocalix[4]pyrrole 34.** Compound 12 ('β-hook' acid) (250 mg, 0.51 mmol) was dissolved in DMF (20 mL, dry) under an argon atmosphere. Methylaminoferrocene (110 mg, 0.51 mmol) was added followed by BOP PF₆ (273 mg, 0.61 mmol) and triethylamine (105 mg, 1.03 mmol). The reaction mixture was stirred for 72 hours. The DMF was then removed *in vacuo* and the product purified by column chromatography (SiO₂, CH₂Cl₂: CH₃OH 99:1 eluant) affording an orange foam **34** (294 mg, 84 %). ¹H NMR (250 MHz, CD₂Cl₂) δ: 8.52 (s, 1H, NH), 7.38 (s, 1H, NH), 7.06 (s, 1H, NH), 7.01 (s, 1H, NH), 6.06 (s, 1H, NH_{py}), 5.91 (s, 6H, CH_{py}), 5.74 (1H, d, CH_{py}), 4.21 (m, 6H, FcH + FcCH₂), 4.14 (s, 5H, 5FcH), 3.54 (s, 2H, OCH₂), 1.58 (m, 12H, CH₃), 1.49 (m, 12H, CH₃). ¹³C NMR (62.90 MHz, CD₂Cl₂) δ: 172.6, 139.9, 139.4, 139.1, 138.9, 138.4, 138.3, 137.6, 134.3, 110.5, 107.1, 103.3, 103.2, 103.1, 102.9, 102.6, 102.1, 85.7, 68.9, 68.4, 68.3, 39.1, 37.1, 35.9, 35.5, 35.2, 29.2, 29.1, 29.0, 28.8. High resolution FABMS calc for C₄₁H₄₉N₅O⁵⁶Fe 683.3287; found 683.3280 (Δ 1.0 ppm)




The electrochemical behavior of these macrocycles includes multiple oxidation processes occurring in the calixpyrrole ring. The ferrocene oxidation wave is distinct from these macrocycle oxidations and can be followed using standard electrochemical techniques (cyclic and square wave) (FIG. 11). The ferrocene oxidation wave is perturbed upon addition of anions to the electrochemical solution (FIG. 12) (Table 1).

Table 1 Redox potentials (mV) vs Ag/AgCl of the ferrocene moiety in compounds 33 and 34 in the absence and presence of anions (perturbation is shown in brackets) from square wave voltammetry

	Compound 33 (mV)	Compound 34 (mV)
Free ligand	453	463
H ₂ PO ₄ ⁻	401 (-52)	430 (-33)
F ⁻	491 (+38)	491 (+28)
Cl ⁻	530 (+77)	410 (-53)


Larger perturbations are observed for the *meso* compound 33 than for the β - compound 34. It has been demonstrated that calixpyrroles can be incorporated into devices (in this case an electrochemical anion sensor) and operate successfully.

meso-Tetrakispolyethyleneglycol-meso-tetramethyl substituted calix[4]pyrrole 35. A polyethylene glycol derivative of calix[4]pyrrole has been synthesized 35.




Reaction of 4-acetylbutyric acid with triethylene glycol monomethyl ether as solvent in the presence of HCl affords a water-soluble triethylene glycol ketone. Pyrrole (48 mg, 0.72 mmol) and the triethylene glycol ketone (200 mg, 0.72 mmol) were dissolved in methanol (5 mL) and methanesulfonic acid (1 drop) added. The reaction was stirred overnight and the solvent then removed *in vacuo* affording the water soluble product 35 in 24 % yield.

Preparation of amidosaccharide-calix[4]pyrrole conjugate 36. The coupling of protected glucosamine with calixpyrrole β -acid 12 derivative of calix[n]pyrrole produces a water soluble calixpyrrole conjugate. Calixpyrrole β -acid 12 (48.67 mg, 0.1 mmol) was dissolved in dry dichloromethane (50 mL) under argon and diisopropylcarbodiimide (2.5 molar equivalents, 0.25 mmol) was added together with 1 mg of 1-hydroxybenzotriazole. The solution was stirred for 10 minutes at room temperature under argon. 2-Amino-2-deoxy-1,3,4,6-tetraacetyl- β -D-glucopyranose hydrobromide salt (2 molar equivalents, 73 mg, 0.2 mmol) was added followed by 0.3 mL of dry pyridine. The reaction mixture was stirred under argon for 3 days, diluted with dichloromethane (50 mL), washed with 3% hydrochloride acid, water and the organic phase dried over sodium sulfate. The product was isolated on short silica gel column with dichloromethane as eluant. Yield of calix[n]pyrrole 36 2-amido-2-deoxy-1,3,4,6-tetra-O-acetyl- β -D-glucopyranose conjugate is 73.4 mg (90%). HR MS for $C_{44}H_{57}N_5O_{10}$ calc. 815.41051, found. 815.41049 Deprotection of tetraacetylated derivative was performed by standard procedure by dissolution of the protected sugar calix[4]pyrrole conjugate described above (50 mg) in MeOH, with catalytic amount of sodium methoxide and stirred for 8 hours at room temperature, the product - calix[4]pyrrole 2-amido-2-deoxy- β -D-glucopyranose 37 conjugate was obtained in quantitative yield. For $C_{36}H_{49}N_5O_6$ calc 647.36826, found 647.36802.



36



37

A host of hydrophilic substituents can be coupled to β -mono-acid and *meso*-mono-acid; for example, polyethylene glycols, azacrown ethers, crown ethers, phosphate groups, saccarides, modified sugars bearing amino groups, amino alcohols, sulfonyl, chitosan, 2-aminoethylhydrogensulfate, 2-aminoethylphosphoric acid, and the like. Additionally, water solublizing groups, such as sulfonates, phosphates, polyalcohols, polyethers, hydroxyl groups, amino groups, carboxylic acid groups, sulfoxide groups, and other could be incorporated into the molecule *via* attachment to the starting ketone and pyrrole before formation of the macrocycle.


Calix[n]pyrroles have been shown to bind phosphate groups in the solid state, in solution, and when bound to solid supports as described in Example 10. DNA and RNA, important biological molecules, contain many phosphate groups. A water-soluble calix[n]pyrrole is expected to bind to DNA and RNA. Binding could have the effect of limiting the ability to unwind in the case of DNA, thereby hindering biological functions, or in the case of RNA, could prevent translation. Additionally, oligonucleotides having sequence complementarity to a template molecule may be attached to a calix[n]pyrrole for site-specificity. Uses include inhibition of replication and anti-sense therapy.

Example 9

**Chiral Calix[n]pyrroles for Recognition and Separation
of Neutral and Anionic Chiral Molecules.**

The present example provides methods for inducing chirality in calix[n]pyrroles. A first method is to use chiral molecules as starting materials for the synthesis of calix[n]pyrroles. A second method is to attach optically active (chiral) groups to the molecule *via* bond formation between functional groups on the calix[n]pyrrole and the chiral auxiliary. A third method is based on transforming achiral substituents into chiral groups by using, for instance, chiral catalysis.

Exemplary chiral ketones that can be used to synthesize chiral calixpyrroles include, but are not limited to, chiral ketone containing steroid derivatives, antibiotics e.g. lasalocid, synthetic chiral ketones, and binaphthal groups that possess an inherent chirality. Steroid analogs have the added advantage of forming bowl-shape cavities because of the natural bend in the molecule. They have well defined geometries and can act as receptors for anions (e.g. 38).




Preparation of steroidal calix[n]pyrroles. 4-Cholesten-3-one (38.4 mg, 0.1 mmol) was dissolved in 25 mL of ethanol. Ten molar equivalents of pyrrole was added to the solution. The solution was cooled to 0° C and 10 mg of concentrated HCl in 1 mL of ethanol was added to the reaction mixture which was then stirred for 24 hrs. at room temperature. The solvent was removed *in vacuo*. The residue was redissolved in dichloromethane (25 mL) and washed with a 3% sodium carbonate solution. The product 38 was purified by flash chromatography on silica gel using a dichloromethane-hexane (1:1) mobile phase. Yield = 12%. The main product was a calix[4]pyrrole (C₃₁H₄₇N)₄ Calc'd Mass: 1734. Found: 1734. For

calix[5]pyrrole ($C_{31}H_{47}N_5$) Calc'd Mass: 2167. Found: 2167. This reaction has been repeated varying the ratio of ketone to pyrrole and also varying the type of acid used. The total yields of calix[n]pyrroles predominantly but not exclusively the $n=4$ product, were generally in the 10-15% range.

5 Because of their many functional groups, steroid-substituted calix[n]pyrroles containing functional groups capable of being attached to solid supports are likely to exhibit enantioselective separation of, for example, sugars, synthetic oligonucleotides, polypeptides, chiral anions, chiral neutral molecules, or the like. One skilled in the art, in the light of the present disclosure, would appreciate that the resulting species would be amenable to
10 attachment to a solid support.


Binaphthal system also possess an inherent chirality. Being so close to the periphery, they could act as "directing groups" and influence binding of neutral and anionic species e.g. 39.



15 A second method for inducing chirality in a molecule is to attach a chiral auxiliary to the main body of the calix[n]pyrrole. This could be done, for example, by forming an amide bond between a carboxy calix[n]pyrrole and a chiral, amino group-containing molecule. This can include, but is not limited to, aminoglycosides, chiral amino alcohols, aminocyclodextrins and amino acids. Many different kinds of functional groups can be coupled to form chemical bonds
20 with a calix[n]pyrrole and would be known in the light of the present disclosure. One skilled in the art, in the light of the present disclosure, would appreciate that the resulting species would be amenable to attachment to a solid support.

Chiral Amide β -Substituted calix[4]pyrrole 40. Monoacid 12 has been coupled to a number of different amines to produce chiral and dimer calixpyrrole species. (R)-(+)- α -

methylbenzylamine is a readily available chiral amine. This amine was appended to the calixpyrrole skeleton by conducting an amide coupling reaction between the amine and compound **12** in the presence of BOP and Et₃N in DMF solution. The amide was isolated as a DMF complex.



Chiral calixpyrroles are useful in the chiral separation of racemic mixtures of species such as amino acids. Compound **40** will form a diastereomeric mixture when complexed to fluoride (FIG. 13).

Compound **12** (200 mg, 0.41 mmol) and (R)-(+)- α -methylbenzylamine (50 mg, 0.41 mmol) were dissolved in dry DMF (10 mL) and stirred under an argon atmosphere. Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP) (0.218 g, 0.494 mmol) and triethylamine (83 mg, 0.82 mmol) in 15 mL DMF were added, and the reaction mixture was stirred overnight. The solvent was then removed under high vacuum and the residue was purified by column chromatography (twice on SiO₂, CH₂Cl₂ - 2.5% MeOH), affording the chiral amide **40** as a DMF complex (161 mg, 66% yield). ¹H NMR (250 MHz, CD₂Cl₂) δ : 8.76 (s, 1H, NH), 7.95 (DMF), 7.33 - 7.28 (m, 5H, PhH), 7.16 (s, 1H, NH), 6.96 (2 br s, 2H, NH), 5.86 - 5.65 (overlapping multiplets, 7H, CH_{py}), 5.09 (m, 1H, PhCH), 3.50 (s, 2H, CCH₂), 2.90 (DMF), 2.81 (DMF), 1.54 - 1.45 (overlapping singlets, 27H, CH₃). High resolution FABMS calc for C₃₈H₄₇N₅O₁ 589.3781; found 589.3768 (Δ 2.2 ppm).

20 The third method of inducing chirality is reaction of a non-chiral substituent, such as an alkene, to form a chiral side chain and subsequent anti-selective transformation of the side chain to a chiral moiety. These types of reactions include, but are not limited to, catalytic asymmetric hydrogenation, asymmetric epoxidation and subsequent ring opening, asymmetric diol formation, asymmetric halogenation and asymmetric amino alcohol formation.

Example 10

Calix[n]pyrroles, Calix[m]pyridino[n]pyrroles and Calix[m]pyridines for Solution and Solid Support Binding and Separation of Anionic and Neutral Substrates

Calixpyrroles possess a pyrrolic hydrogen bonding array capable of coordinating anions *via* N-H...anion hydrogen bonds. The anion binding properties of calixpyrroles have been studied in solution using ^1H NMR titration techniques and are provided herein.

Stability constants for octamethylcalix[4]pyrrole, **1**, and tetraspirocyclohexylcalix[4]pyrrole, **2**, binding to various anions in dichloromethane were obtained and are provided in Table 2.

Table 2: Stability constants for compounds **1 and **2** with anionic substrates^a**

		Stability Constant (M^{-1})	
		Calixpyrrole 1	Calixpyrrole 2
	Fluoride ^b	17170 (± 900)	3600 (± 395)
15	Chloride	350 (± 5.5)	117 (± 4.0)
	Bromide	10 (± 0.5)	^c
	Iodide	< 10	^c
	Dihydrogen phosphate	97 (± 3.9)	< 10
	Hydrogen sulfate	<10	^c

^aAnions were added as 0.1M CD_2Cl_2 solutions of their tetrabutylammonium salts to receptor to 10 mM receptor solutions in CD_2Cl_2 with concentration changes being accounted for by EQNMR. ^bTetrabutylammonium fluoride was added as the trihydrate. ^cUndetermined.

The data of Table 2 demonstrate that calix[4]pyrroles show different affinity for different anions in solution, with fluoride being the most tightly bound while iodide is very slightly bound.

The present inventors have discovered that varying the substituents on the β -position of a calixpyrrole molecule, however, affects the strength with which a particular anion is held. The data of Table 3 shows the effect of substituting the β -positions of the macrocycle with ester **10**, methoxy **13**, and bromine **14** groups and the effect that these functional groups have on the binding strength of calix[4]pyrrole to various anions. Additionally compound **11** has been shown to interact with both fluoride and chloride anions in solution.

Table 3: Stability constants for compounds 10, 13 and 14 with anionic substrates^a in CD₂Cl₂

		<u>Stability Constant (M⁻¹)</u>		
		10	13	14
5	Fluoride	1.1 (± 0.2) × 10 ³	1.7 (± 0.2) × 10 ²	2.7 (± 0.4) × 10 ^{4b}
	Chloride	47 (±1)	<10	4.3 (±0.6) × 10 ³
	Dibiphosphate	^c	6.5 (± 0.4) × 10 ²	<10
10	^a Anions were added as 0.03 M CD ₂ Cl ₂ solutions of their tetrabutylammonium salts to 3 mM solutions of the receptor in CD ₂ Cl ₂ with concentration changes being accounted for by EQNMR. In determining the stability constants, the possible effects of ion pairing (if any) were ignored.			
	^b Estimated value. The NH proton resonance broadened considerably during the titration, forcing the frequency of the resonance to be noted manually.			
	^c Not determined.			

15

The attachment of electron-donating substituents (such as methoxy) to the β-position (carbon or C-rim of the calixpyrrole) decreases the stability constants of compound 13 relative to compound 2. However attachment of electron-withdrawing substituents (such as bromine) to the carbon-rim has increased the stability constants of compound 14 relative to compound 1.

20 While varying the electron density of the calix[4]pyrrole core increases or decreases the binding affinities towards anions, the present inventors also discovered that groups having a functional moiety at a suitable distance from the macrocycle, and having sufficient flexibility to fold back over the macrocycle also affects the strength of binding as discussed below. The data of Table 4 illustrate the effects of an amide group attached at the *meso*-position of the

25 calix[4]pyrrole.

Table 4: Stability constants for compounds 28 and 29 with anionic substrates in CD₂Cl₂

		<u>Stability Constants (M⁻¹)</u>	
		Compound 28 (β)	Compound 29
30	(<i>meso</i>)		
	Chloride	405 (± 10)	415 (± 45)
	Dihydrogen phosphate	80 (± 15) ^b	62 (± 6)
	Hydrogen sulfate	< 10	< 10

35 ^aAnions were added as 0.1 M CD₂Cl₂ solutions of their tetrabutylammonium salts to 10 mM solutions of the receptor in CD₂Cl₂ with concentration changes being accounted for by EQNMR. In determining the stability constants, the possible effects of ion pairing (if any) were ignored.

40 ^bEstimated value. The NH proton resonance broadened considerably during the titration, forcing the frequency of the resonance to be noted manually.

As compared to tetraspirocyclohexylcalix[4]pyrrole, **2**, the *meso*-substituted, amide-substituted calix[4]pyrrole **29** shows a marked increase in binding affinities for chloride and phosphate when compared with the values obtained for tetraspirocyclohexyl calix[4]pyrrole (see Table 2). The amide group contributes to the binding of the anion *via* an amide-anion hydrogen bond as shown in FIG. 14.

This increased binding affinity towards anions was not observed for the β -substituted amide substituted *meso*-octamethylcalix[4]pyrrole **28** when compared to the compound **1**. A possible reason is that the carboxy "tail" is not long enough to swing around and contribute to the binding of the anion.

10

Table 5 Stability constants of the ferrocene-substituted calix[4]pyrroles **33 and **34**, measured in CD_2Cl_2 .**

Anion	33 <i>meso</i> -ferrocene	34 β -ferrocene
H_2PO_4	$40 \pm 6 \text{ M}^{-1}$ (β -CH)	$40 \pm 9 \text{ M}^{-1}$
15 F	$79 \pm 7 \text{ M}^{-1}$ (β -CH) $206 \pm 51 \text{ M}^{-1}$ (NH)	$1496 \pm 370 \text{ M}^{-1}$
Cl	$202 \pm 15 \text{ M}^{-1}$	$444 \pm 22 \text{ M}^{-1}$

The ferrocene modified calixpyrrole binding constants with anions are provided in Table 5.

20 **Neutral Substrates Binding in Solution.** The binding of neutral molecules to calix[4]pyrrole in the solid state prompted the present inventors to quantify the binding affinities of calix[4]pyrrole to a number of hydrogen-binding neutral molecules. Table 6 illustrates the varying degrees of affinity to which calix[4]pyrrole binds to various hydrogen binding compounds.




Table 6: Association Constants for Calix[4]pyrrole, 1, with Neutral Substrates^a

	Substrate Added	K_a (M^{-1})
	methanol	12.7 ± 1.0
	ethanol	10.7 ± 0.7
5	benzyl alcohol	9.7 ± 0.7
	isopropyl alcohol	7.0 ± 0.4
	<i>sec</i> -butanol	6.2 ± 0.4
	<i>N</i> -formylglycine ethyl ester	13.3 ± 1.0
	<i>N,N</i> -dimethylformamide	11.3 ± 0.8
10	<i>N,N</i> -dimethylacetamide	9.0 ± 0.9
	1,1,3,3-tetramethylurea	2.2 ± 0.1
	dimethyl sulfoxide	16.2 ± 1.1
	1,2-dimethylimidazole	5.4 ± 0.3
	acetone	2.2 ± 0.2
15	nitromethane ^b	—

^a In benzene-*d*₆ at 298 K. For each titration, the concentration of **1** was held constant (at *ca.* 4×10^{-3} M) as aliquots of the substrate in benzene-*d*₆ (*ca.* 1 M) were added. ^b In this instance, the induced shifts in the NH proton(s) of **1** were too small (< 0.15 ppm) to afford a reliable K_a value.

20 The data in Table 6 demonstrates the ability of calixpyrroles to coordinate to neutral species

Molecule Separation using Solid-Supported Calix[4]pyrroles. Following the discovery of the unique anion binding capabilities of calix[4]pyrroles, the present inventors proceeded to attach the “monohook” of Examples 2 and 3 to a silica gel solid support. 25 Calix[4]pyrrole-substituted silica gel chromatography has demonstrated the ability to separate anionic, neutral hydrogen-binding and aromatic molecular species. In addition to anionic species such as phenyl phosphate, calix[4]pyrrole substituted silica gel has demonstrated the ability to separate polyphosphate-containing compounds such as adenosine mono-, di-, and triphosphate under standard, isocratic HPLC conditions at pH = 7. As one might expect, increasing the 30 number of anionic binding sights greatly increases the retention times of these types of molecules.

The present approach to anionic and neutral species separation was as follows. The macrocycle, *meso*-tris-spirocyclohexyl-monohook-calix[4]pyrrole, **4**, was covalently attached to 5 μ , trimethyl silyl protected, aminopropyl HPLC-grade silica gel obtained from Phase 35 Separations (Norwalk, CT) to produce Gel M as shown in FIG. 15A. In a similar fashion, β -monohook acid calix[4]pyrrole **12** was attached to 5 μ , trimethyl silyl protected, aminopropyl HPLC-grade silica gel obtained from Phase Separations (Norwalk, CT) in the presence of diisopropylcarbodiimide to produce Gel B as shown in FIG. 15B. The attachment of the

"monohook" calix[4]pyrrole to C-1 aminopropyl silica gel is achieved in a greater than 95% yield.

The calix[4]pyrrole columns were packed commercially by Alltech (Deerfield, IL), and various substrates, under differing conditions, were eluted through the columns. All of the data provided in the chromatograms herein were obtained at room temperature under isocratic conditions. FIG 16 illustrates the separation of phenyl sulfonate (3), phosphate (4), arsenate (1), and benzoic acid (2) on Gel M. Excellent selectivity is seen among the different anions. FIG. 17 and FIG. 18 provide further examples of separation of biologically-relevant substrates by the calix[4]pyrrole-substituted silica gel, in this case, adenosine monophosphate, adenosine diphosphate and adenosine triphosphate. The nucleotides described elute in ascending order according to the number of phosphate groups contained within the molecule (i.e. the substrates containing the more anionic groups are retained longer on the column relative to substrates containing fewer anionic groups). Selectivity for carboxylates was also observed for carboxy-substituted benzenes. In two separate experiments, calix[4]pyrrole substituted silica gel, B, effectively separated benzoic acid, isophthallic acid, and 1,3,5-tricarboxy benzene. In a separate experiment, a calix[4]pyrrole substituted silica gel column effectively separated phthallic acid from a mixture of teraphthallic acid, isophthallic acid and phthallic acid.

FIG. 19 and FIG. 20 demonstrate the separation of carboxybenzyl-N-protected amino acids on Gel M and Gel B, respectively. These data illustrate that both hydrophobic contributions and hydrogen bonding interactions are involved in the separation.

FIG. 21 illustrates the separation observed between 5 different neutral aromatic compounds: uridine (1), phenol (2), aniline (3), benzene (4), and 4-iodo-nitro benzene (5) on Gel M. In addition to hydrogen-bonding properties, π - π interactions appear to influence separation. It is noteworthy to point out that nucleotides, shown in FIG. 17 and 18, require a substantially stronger buffer system for elution than the nucleoside, uridine (1), shown in FIG. 21, which requires a weak buffer and is eluted almost immediately.

Further, the separation of very similar molecules of a class of molecules was carried out. Using a model system analogous to the class of polychlorobiphenyls which are known carcinogens, the separation of polyfluorobiphenyls, a biologically inactive compound, was effected. Demonstrated in FIG. 22, is the separation of four polyfluorobiphenyl compounds; 4-fluorobiphenyl (1), 4,4'-difluorobiphenyl (2), 2,2',3,3',5,5',6,6'-octafluorobiphenyl (3), and perfluorobiphenyl (4) on Gel M and in FIG. 23 on Gel B.

With current interest in antisense oligonucleotide therapy and oligonucleotides as primers and probe molecules, the ability to effectively and efficiently purify usable quantities of these oligomers is of paramount importance. One of the best methods for obtaining large quantities of purified oligonucleotides is high pressure liquid chromatography (HPLC). There are a number of HPLC techniques employed in the purification of oligonucleotides, however, the two most currently used methods are reverse phase- and ion exchange chromatography.

These popular methods suffer from practical drawbacks. Purification of medium chain length (12-24) oligonucleotides using reverse phase chromatography must be done using protected substrates. The protecting groups most often used are trityl groups which must be subsequently removed by hydrolysis before use. Additionally, reverse phase chromatography suffers from very long retention times which limits its applicability. Oligonucleotides separated *via* anion exchange chromatography, while not plagued by the long retention times or additional hydrolysis steps, must be desalted before use.

FIG. 24 illustrates a separation of 12 to 18 base mixture of deoxythymidilates (dT₁₂₋₁₈) on Gel M and FIG. 25 shows the data for Gel B. Additionally mixtures of deoxythymidilates (dT₁₉₋₂₄) were successfully separated on both Gel M and Gel B. The calix[4]pyrrole-modified silica gel has some immediate advantages over anion exchange columns and reverse phase columns. With respect to anion exchange chromatography, the large concentrations of salt that is normally need to effect separation is not needed, thereby removing the mandatory desalting step after chromatography of oligonucleotides before using them in biochemical experiments. The primary advantage that the calix[4]pyrroles has over reverse phase columns is the higher resolution and shorter retention times. Additionally, reverse phase chromatography of oligonucleotides is done while they are in a "protected" state and must further be hydrolyzed before use.

The ability of calix[4]pyrroles to distinguish between specific sequences of nucleotide hexamers was then examined. FIG. 26 provides the data where three different oligonucleotide hexamers, GGGCCC, TCTAGA, and GCATGC, were eluted through Gel M resulting in separation of the three hexamers.

Mixtures of pesticides (carbofuran, carbendazim, bromacil, bentazon, carboxin and norflurazon) were successfully separated on Gel M.

Oligonucleotide therapy has been proposed for a wide range of ailments. These include, but are not limited to sickle cell anemia, hypertension, leukemia, cancer, and AIDS. The basic premise of oligonucleotide therapy is to identify the relevant mRNA or DNA strand that is causing the specific problem and synthesizing a short (≤ 20 bases) complementary oligonucleotide strand that will bind to the mRNA or DNA, thereby blocking translation or transcription of the genetic material.

Table 7 illustrates the ability of calix[4]pyrrole silica gels M and B to separate tetrabutyl ammonium halides.

TABLE 7: Analysis of Tetrabutylammonium Halides on Calixpyrrole-Substituted Silica Gel

		Elution Times (minutes)	
		Silica Gel B	SilicaGel M
5	Chloride ^b	17.9 (± 0.1) ^a	15.2 (± 0.1) ^a
	Dihydrogen phosphate ^b	22.0 (± 0.1) ^a	20.1 (± 0.1) ^a
	Hydrogen sulfate ^b	16.2 (± 0.1) ^a	16.2 (± 0.2) ^a
	Fluoride ^b	16.9 (± 0.1) ^a	16.4 (± 0.2) ^a

10 ^aTimes given for individual elution of tetrabutyl ammonium anions. Separation of mixtures was not possible due to broadening of peaks. ^bAnions were eluted as 1 mM CH₃CN solutions of their tetrabutylammonium salts under the following conditions: Mobile phase, CH₃CN; Flow Rate, 0.40 mL/min.; Detection, Conductivity; Column Temperature, 25° C.

15 The present example primarily refers to silica-bound calixpyrrole, however, the solid support attached to the calix[4]pyrrole is not limited to silica gel. Further solid supports include, but are not limited to, polystyrene, polyacrylamide, Merrifield resins, glass, sepharose, sephadex, agarose, clays, zeolites, and the like, that can be functionalized to allow the formation of a calix[n]pyrrole bonded to a solid support.

20 Calix[n]pyrroles have been shown to be selective toward a number of different anionic species. In examples described herein, calix[4]pyrrole has shown differing affinities for anionic species including fluoride, chloride, bromide, iodide, phosphates, carboxylates, arsenates and sulfates.

25 Macrocycles of the present invention may be used to generate ion selective electrodes in a further use embodiment. Data illustrated in Table 8 for anion selective electrodes synthesized by the present inventors indicate that the selectivity and preferential binding observed in solution and in separation chemistry, described in examples herein, is consistent with anion selective electrode technology.

Table 8: Anion Selectivity for a Calix[4]pyrrole Based Membrane^{a,b}

		$\log K^{\text{pot}}_{ij}$
	Fluoride	0
	Chloride	-0.7
5	Bromide	-1.0
	Iodide	+0.4
	Hydrogen Sulfate	-4.4
	Dihydrogen Phosphate	-1.7
	Hydrogen Sulfite	-0.4
10	Nitrate	-0.5
	Nitrite	-1.8
	Acetate	-0.9
	Benzoate	+0.02
	Benzene sulfonate	-0.5
15	L-Alanine	-1.7
	L-Aspartate	-1.6
	L-Tryptophan	-0.5
	L-Hystidine	-1.6
	D-Serine	-1.1
20	DL-Alanine	-1.2
	DL-Threonine	-1.4
	Chromotropate	-1.0
	5-Formyl-2-furansulfonate	-0.6
25	^a Selectivity coefficients determined relative to fluoride, $\log K^{\text{pot}}_{ij}$, using the separate solution method. ^b Membrane compositions: 3% (w/w) calix[4]pyrrole, 22% (w/w) PVC, 75% (w/w) 2-onitrophenyloctyl ether. Internal electrolyte was 0.1 M NaCl and 0.001 M NaF.	

Methods of making calix[n]pyrrole-ion-selective electrodes include, but are not limited to, the following examples: attaching, or immobilizing a functionalized calix[n]pyrrole to a functionalized polymer and further coating this polymer to an electrode that is sensitive to changes in ionic strength; immobilizing the calix[n]pyrrole in a plasticizers and encasing the mixture in a membrane surrounding an electrode that is sensitive to changes in ionic strength; covalently attaching the calix[n]pyrroles to an electrode that is sensitive to changes in ionic strength; or coating the electrode with a layer of calix[n]pyrroles containing electropolymerizable functional groups, including but not limited to, pyrrole, thiophene or vinyl, and polymerizing the coating on to the electrode.

Example 11

Biomedical applications

The anion and neutral binding properties of calix[n]pyrrole, calix[m]pyridino[n]pyrrole and calix[m]pyridines allow for a number of useful applications within the biomedical field.

5 Applications include, but are not limited to, drug-delivery systems, anion transport through membranes, selective anion channel formation, dialysis, blood filtration and viral inhibition.

Cystic fibrosis is the most common lethal genetic disease among Caucasians, affecting one out of 2,500 infants in the U.S. The disease is characterized by the inability to produce properly-functioning chloride anion channels. The calix[n]pyrroles have been demonstrated
10 herein to bind chloride *via* hydrogen bonding. An array of calix[n]pyrroles or calix[m]pyridino[n]pyrroles could be designed to span a membrane requiring selective chloride transport. Chloride anion is bound and released appropriately by calix[4]pyrrole as demonstrated in Example 10. Calix[4]pyrrole has also demonstrated the ability to transport anions through an "artificial" dichloromethane membrane in Pressman U-tube experiments. An
15 array of calix[n]pyrroles is expected to transfer chloride anions by "handing off" the anion to the next calix[n]pyrrole.


Calix[n]pyrroles can also be used to transport the anion through the membrane by acting as discrete carriers. Referring to the U-tube data, calix[4]pyrrole has demonstrated the ability to transfer charged molecules through organic membranes. The chloride anion could be transferred
20 through a cell membrane in this way thereby alleviating the problem of chloride transport and ultimately treating the disorder.

Another application of anion transport is use of the macrocycles of the present invention as a drug delivery system. Many potential anti-viral agents which display activity *ex-vivo* are phosphorylated nucleosides. These species are too polar to pass through cell wall
25 membranes and are thus inactive *in vivo*.

Calix[4]pyrrole has been shown to transport anionic species effectively through "artificial" membranes. The present inventors have synthesized a number of nucleotide conjugates as examples of nucleobase-substituted calix[n]pyrroles. The possibilities of nucleotide substitution include, but are not limited to any of the naturally occurring purine or
30 pyrimidine nucleobases. Additionally, modified versions include any number of purine- or pyrimidine- based synthetic molecules capable of addition to a calix[4]pyrrole core. "Nucleobase," as used herein, means a purine or pyrimidine base, nucleoside (saccharide derivative), nucleotide (saccharide-poly- or mono-phosphate derivative) and natural and synthetic analogues thereof. Attachment of a nucleobase to the calix[n]pyrrole may occur at a
35 functional group of the β -position, a pyrrolic-nitrogen, or *meso*-position of the calix[n]pyrrole. One nucleobase may be attached, however, any number or diversity of nucleobases may be attached to the calix[n]pyrrole. Attachment can occur using methods such as: amide bond formation, Stille coupling, ester formation, or the like. The nucleobase appended calixpyrrole

will coordinate to both the phosphate group and nucleoside of the anti-viral drug, forming a hydrophobic complex which may pass through cell wall membranes and decomplex, releasing the drug into the interior of the cell.

Examples of nucleobase-calix[4]pyrrole conjugates synthesized by the present inventors are provided as macrocycles 41 and 42.



Preparation of nucleobase substituted calix[n]pyrroles 41 and 42 (general procedure).

Calixpyrrole monoacid (1 mmol) was dissolved in dry dichloromethane (25 mL), (chloroform, acetonitrile, DMF, 1,2-dichloroethane can also be used). The solution was activated with 2 molar equivalents (per one acid group) of the activating agent diisopropylcarbodiimide at room temperature in the presence of five molar percent of 1-hydroxybenzotriazole and 4-dimethylaminopyridine. Two molar equivalents per carboxy unit of an N-protected amino acid were added. The reaction mixture was stirred at room temperature for 24-36 hours. Acetic acid (0.5 mL) was added, followed by multiple washings with a 3% hydrochloric acid solution, water, a 5% solution of sodium hydrogencarbonate and water. After drying over sodium sulfate the product was obtained by crystallization or flash chromatography on silica gel. Removal of the N-protecting group was done using standard procedures. The removal of trityl- and benzyl-protecting groups were achieved using trifluoroacetic acid for trityl and ammonia in methanol for benzyl protecting groups.

Studies have demonstrated that these nucleobase-calix[4]pyrrole conjugates are not only efficient carriers, but selective carriers as well. Using a cytosine-calix[4]pyrrole conjugate to transport a mixture of adenosine monophosphate, cytidine monophosphate, guanine monophosphate, thymidine monophosphate and uridine monophosphate, a transfer ratio of 5:1:125:8:7. The guanine monophosphate was transported from one aqueous phase to another 10-100 times better across the dichloromethane model membrane as a result of having complementary binding with the cytosine of the cytosine-calix[4]pyrrole conjugate. In addition to being bound to a single nucleobase, calix[n]pyrroles could be attached to oligonucleotides using known technology. Calix[n]pyrrole-oligonucleotide conjugates could then be used to bind such substrates as mRNA or DNA preventing transcription or translation. This mode of binding could be used in antisense therapy, or viral inhibition. In the case of viral inhibition, the binding site of the virus would be identified and the complementary calix[n]pyrrole-oligonucleotide conjugates would be synthesized.

Calix[n]pyrroles have utility for *ex vivo* applications as well. Kidney failure is a condition that affects a significant portion of the people in the world due to various ailments. The present example provides for the use of calix[n]pyrroles and calix[m]pyridino[n]pyrroles for dialysis or filtration of *ex vivo* bodily fluids. Phosphates, chloride, and other toxins contained in the blood or other bodily fluids may be removed by exposure to macrocycles of the present invention since said macrocycles have demonstrated the ability to bind anionic and neutral molecules in the solid state, in solution and when attached to a solid support. Further, calix[n]pyrroles can be functionalized to be anion-specific. The present inventors have shown that under chromatographic conditions where anions are bound, proteins and other biomolecules were unhindered in passage through a column, for example.

Calix[n]pyrroles could also be used in dialysis. Current dialysis employs an anion gradient to bring anions across a membrane, thereby establishing an equilibrium. Calix[n]pyrroles could be used to bind the anions once they came across the membrane forcing the equilibrium to shift in favor of additional waste, anion removal by not allowing an equilibrium to form. The calix[n]pyrroles could either be water soluble and partitioned on one side of a membrane or bound to a solid support to prevent diffusion into the blood supply.

Calix[n]pyrrole could be attached to a biologically-inert solid support and used to filter toxins out of the blood. These toxins include, but are not limited to, drug metabolites (for use in drug overdoses) and anionic wastes. Another endogenous material that can function as a toxin is urea. Its removal using this approach illustrated could be used in the treatment of gout.

Additionally calixpyrroles could be used in imaging. ^{99m}Tc is a high energy form of technetium used in imaging for the detection of tumors. $^{99m}\text{TcO}_4^-$ or any other anion containing ^{99m}Tc may be coordinated to a calix[n]pyrrole and used as an imaging agent.

Example 12**Environmental Remediation**

Radioactive waste poses a serious problem to the environment. Current technology generates large amounts of secondary solid waste for disposal. Separation and concentration of these wastes is of paramount importance, especially due to long half-lives, and inadequate methods of storage such as sludges or underground tanks of wastewater. Half-life examples for Tc-99, Ni-63, Cs-137, and Sr-90 are 213,000 years, 100 years, 30 years and 28 years, respectively.

Certain of the radioactive metallic wastes exist in an oxidized anionic form and are therefore soluble in aqueous solutions. For example, technetium exists as TcO_4^- and nickel, strontium, and cesium can exist in oxidized anionic forms.

Calix[4]pyrrole has demonstrated the ability to bind phenyl arsenate in aqueous media. In addition, FIG. 16 illustrates the ability of calix[4]pyrrole to separate 4 different phenyl anions from a mixture thereof.

In a column embodiment, solid-supported calix[n]pyrrole could be used to separate out metallic anions from a stream passing through a column. The bound anions could then be eluted from the solid support, and the concentrated eluate is able to be stored in a smaller volume.

In a batch embodiment, solid support-bound calix[n]pyrrole could be mixed with a waste solution containing an anion to be removed and the mixture then filtered as a method of extracting the waste anions out of solution. Additionally, a multi-phase extraction system is envisioned where calix[n]pyrrole in a first phase extracts anionic waste from another phase, and the waste then is separated for storage or disposal.

The present inventors envision, because of the relatively fast "on-off" rate that is observed with calix[4]pyrrole, that the radioactive waste metal could be "washed" off of the solid support for solid support reuse. Current technology uses materials with high binding constants for removal of metallic oxyanion waste. This procedure is efficient for removal of the waste from contaminated liquid, however, the waste is converted to solid waste and must still be disposed of.

Molecules that can be removed from an environmental source are those ions and molecular species that calix[n]pyrrole will bind as described herein. In particular, for application to inorganic metal oxoanions, arsenate, tungstenate, pertechnetate, borate, or the like, may be removed. Removal of pertechnetate from nuclear waste is a particularly preferred application of the present technology. Types of solid supports for attaching calix[n]pyrrole include, but are not limited to, those supports provided in Example 10.

Eutrophication is a serious problem for bodies of water near agricultural lands. The accumulation of phosphates and nitrates from fertilizers in lakes, rivers and inland waterways causes toxic algal blooms and chokes the water of life. Algal blooms are also thought to be toxic for humans. Removal of such anions by use of macrocycles of the present invention in a

water-treatment plant, for example, would purify the water and provide it for human use and generations of the future.

Purification of domestic water supplies is an increasing necessity in densely populated areas. The macrocycles of the present invention would be useful in filters for attachment to
5 home water supplies for removal of anionic pollutants such as, but not limited to, fluoride, phosphate and nitrate.


All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the
10 compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More
15 specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

20



WE CLAIM:

1. A calix[n]pyrrole macrocycle noncovalently-complexed to a molecular or anionic species where n is 4, 5, 6, 7, or 8.
2. A calix[n]pyrrole macrocycle attached to a solid support where n is 4, 5, 6, 7, or 8.
3. A calix[n]pyrrole macrocycle having a first conformation in a solid state when unbound to a molecular or anionic species and having a second in a solid state when bound to a molecular or anionic species where n is 4, 5, 6, 7, or 8.
4. The macrocycle of claim 1, 2, or 3 wherein the calix[n]pyrrole has structure I.



wherein

- 15 when n is 4, p = q = r = s = 0, R₁ - R₁₆ are independently substituents as listed in paragraph i) below, and R_A - R_D are independently substituents as listed in paragraph ii) below;

- when n is 5, $p = 1$, $q = r = s = 0$, R_1 to R_{20} are independently substituents as listed in paragraph i) below, and $R_A - R_E$ are independently substituents as listed in paragraph ii) below;
- 5 when n is 6, $p = q = 1$, $r = s = 0$, R_1 to R_{24} are independently substituents as listed in paragraph i) below, and $R_A - R_F$ are independently substituents as listed in paragraph ii) below;
- when n is 7, $p = q = r = 1$, $s = 0$, R_1 to R_{28} are independently substituents as listed in paragraph i) below, and $R_A - R_G$ are independently substituents as listed in paragraph ii) below;
- 10 when n is 8, $p = q = r = s = 1$, R_1 to R_{32} are independently substituents as listed in paragraph i) below, and $R_A - R_H$ are independently substituents as listed in paragraph ii) below;
- i) hydrogen, halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran, tetrahydrothiapyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;
- 15
- 20
- 25 ii) hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidealkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl; and
- or
- 30 at least two substituents are coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein in paragraph i) or ii) other than for bridged substituents;
- wherein odd-numbered R-substituents are other than hydrogen.
- 35

5. A calix[n]pyrrole macrocycle having structure I:

when n is 8, $p=q=r=s=1$, R_1 to R_{32} are independently substituents as listed in paragraph i) below, and $R_A - R_H$ are independently substituents as listed in paragraph ii) below;

5 i) hydrogen, halide, hydroxyl, alkyl other than methyl or ethyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamido, carboxyamidoalkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, 10 tetrahydropyran, tetrahydrothiapyran, thioalkyl, haloalkyl other than chloropropyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;

15

ii) hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidoalkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl; and

20

or


at least two substituents may be coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein in paragraph i) or ii) other than for bridged substituents;

wherein

25

odd numbered R-substituents are other than hydrogen; and the calix[n]pyrrole is other than *meso*-tetramethyl-*meso*-tetraphenylcalix[4]pyrrole, *meso*-octacyanopropylcalix[4]pyrrole, octaethyl-xanthoporphinogen, 5, 10, 15, 20-*meso*-tetrakis(3,5-di-*tert*-butyl-4-quinomethide)porphyrinogen, or *meso*-octa-3-chloropropylcalix[4]pyrrole.

6. A calix[m]pyridino[n]pyrrole macrocycle of structure II



wherein m designates a number of pyridines in the macrocycle and n designates a number of pyrroles in the macrocycle;

$m+n=4$;

m is other than 1 or 2;

when m is 4, n = 0, p = 1, q = 0, R101 to R108 and R201 to R212 are independently substituents as listed in paragraph i) below, and R301 - R304 are independently substituents as listed in paragraph ii) below;

when m is 3, n = 1, p = 0, q = 1, R101 to R108 and R201 to R211 are independently substituents as listed in paragraph i) below, and R301 - R304 are independently substituents as listed in paragraph ii) below;

i) hydrogen, halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran, tetrahydrothiapyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-

directing molecule, to a catalytic group, to a reporter group, or to a binding agent;

- 5 ii) a lone pair of electrons, hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidealkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl; and

or

- 10 at least two substituents are coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein in paragraph i) or ii) other than for bridged substituents;

wherein R₁₀₁-R₁₀₈ are other than hydrogen;

15

wherein when R₃₀₁-R₃₀₄ is other than a lone pair of electrons, Z is 1;

wherein when R₃₀₁-R₃₀₄ is a lone pair of electrons, Z is 0.

- 20 7. A calix[m]pyridino[n]pyrrole macrocycle
where

m and n designate a number of pyridines and pyrroles in the macrocycle, respectively, m and n are other than 0, and m+n=5, 6, 7, or 8;

- each pyridine or pyrrole α -carbon is bound to another pyridine or pyrrole α -carbon via one non hydrogen-linked sp³ hybridized *meso* carbon;

- 25 each sp³ hybridized *meso* carbon is further independently bonded to a halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl or alkyl ester group; to a site-directing molecule; to a catalytic group; to a reporter group; to a binding agent; or to a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;

35

each pyridine β carbon, pyrrole β carbon and pyridine γ carbon is independently bonded to a hydrogen, halide, hydroxyl, alkyl,

5

alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester group; to a site-directing molecule; to a catalytic group; to a reporter group; to a binding agent; or to a couple that is coupled to a site-directing molecule; to a catalytic group; to a reporter group, or to a binding agent;

10

15


each pyridine or pyrrole nitrogen is bound to a lone pair of electrons, hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidealkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl; and


or

20

25

at least one sp^3 hybridized *meso* carbon, pyridine β -carbon, pyrrole β -carbon, pyridine γ carbon, pyrrole nitrogen or pyridine nitrogen is coupled to form a bridged structure to itself or to another sp^3 hybridized *meso* carbon, pyridine β -carbon, pyrrole β -carbon, pyridine γ carbon, or pyrrole nitrogen; and when coupled to form a bridged structure, non-bridged atoms are as defined for an sp^3 hybridized *meso* carbon, pyridine β -carbon, pyrrole β -carbon, pyridine γ carbon, pyrrole nitrogen, or pyridine nitrogen.





when m is 4, $p = q = r = s = 0$, R₁₀₁ to R₁₁₂ and R₂₀₁ to R₂₀₈ are independently substituents as listed in paragraph i) below, and R₃₀₁ - R₃₀₄ are independently substituents as listed in paragraph ii) below;

when m is 5, p =1, q = r = s =0, R₁₀₁ to R₁₁₅ and R₂₀₁ to R₂₁₀ are independently substituents as listed in paragraph i) below, and R₃₀₁ - R₃₀₅ are independently substituents as listed in paragraph ii) below;

when m is 6, $p = q = 1$, $r = s = 0$, R_{101} to R_{118} and R_{201} to R_{212} are independently substituents as listed in paragraph i) below, and R_{301} - R_{306} are independently substituents as listed in paragraph ii) below;

when m is 7, p=q=r= 1, s =0, R101 to R121 and R201 to R214 are independently substituents as listed in paragraph i) below, and R301 - R307 are independently substituents as listed in paragraph ii) below;

when m is 8, $p = q = r = s = 1$, R₁₀₁ to R₁₂₄ and R₂₀₁ to R₂₁₆ are independently substituents as listed in paragraph i) below, and R₃₀₁ - R₃₀₈ are independently substituents as listed in paragraph ii) below;

- 5 i) hydrogen, halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, 10 alkyl sulfide, tetrahydrotetrapyrane, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;
- 15 ii) a lone pair of electrons, hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidealkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl; and

or

- 20 at least two substituents are coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein in paragraph i) or ii) other than for bridged substituents;

25

wherein R₂₀₁-R₂₁₆ are other than hydrogen;

wherein when R₃₀₁-R_{30m} is other than a lone pair of electrons, Z is 1;

30

wherein when R₃₀₁-R_{30m} is a lone pair of electrons, Z is 0.

9. The macrocycle of any one of claims 5-8 wherein the site-directing molecule is an oligonucleotide, an antibody, or a peptide having affinity for a biological receptor.
- 35 10. The macrocycle of any one of claims 5-8 wherein the binding agent is a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, a calixarene, a cation-binding functionality, a crown ether, a chelating group, a porphyrin, or an expanded porphyrin.

11. The macrocycle of any one of claims 5-8 wherein the reporter group is a redox active group, or a fluorescent molecule.
12. The macrocycle of claim 11 wherein the redox active group is ferrocene.
13. The macrocycle of claim 5, 6, 7, or 8 wherein at least one R substituent attached to a *meso* carbon is carboxy or carboxyalkyl.
14. The macrocycle of claim 5, 6, 7, or 8 wherein at least one R substituent attached to a β - or γ - carbon is other than hydrogen.
15. The macrocycle of claim 5, 6, 7, or 8 wherein at least one R substituent attached to a β - or γ - carbon is carboxy, carboxyalkyl, ester, or carboxamide.
16. The macrocycle of claim 5, 6, 7, or 8 wherein at least one pyrrole or pyridine nitrogen is bound to hydrogen.
17. The macrocycle of claim 5 where n is 4, R_A - R_D are each hydrogen, odd-numbered R groups are methyl, one even-numbered R group is $(CH_2)_tCOOH$ where $1 \leq t \leq 10$, and even-numbered R groups that are other than $(CH_2)_tCOOH$ are hydrogen.
18. The macrocycle of claim 5 where n is 4, R_A - R_D are each hydrogen, each of three *meso* carbons is bridged to itself in a spirocyclohexyl substituent, one *meso* carbon is bonded to a methyl group and to $-CH_2CH_2CH_2COOH$, and even-numbered R groups are hydrogen.
19. A calix[n]pyrrole where n is 5, 6, 7, or 8.
20. A chiral calix[n]pyrrole where n is 4, 5, 6, 7 or 8.
21. A method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having different substituents at *meso*-carbon atoms, comprising
 reacting pyrrole and at least two ketone molecules to produce a calix[n]pyrrole having different substituents at the *meso*-carbon atoms.
22. The method of claim 21 where the number of different ketone molecules includes acetone and cyclohexanone.

23. A method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having an ester substituent, comprising
reacting an ester-functionalized ketone, pyrrole, and a ketone to produce a calix[n]pyrrole having an ester substituent.
- 5
24. The method of claim 23 wherein the ester-functionalized ketone is methyl-4-acetylbutyrate, and the ketone is cyclohexanone, acetone, or pentan-3-one.
25. A method of making a calix[n]pyrrole where n is 4, 5, 6, 7, or 8, the calix[n]pyrrole having an acid substituent, comprising
10 reacting an ester-functionalized ketone, pyrrole and a ketone to produce a calix[n]pyrrole ester derivative; and
de-esterifying the ester derivative to produce a calix[n]pyrrole monoacid derivative.
- 15
26. A method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising activating a calix[n]pyrrole where n is 4, 5, 6, 7, or 8; and
reacting the activated calix[n]pyrrole with an electrophile.
27. The method of claim 26 where the activating is deprotonating the calix[n]pyrrole using a
20 Brønsted base, or a Brønsted base in conjunction with a complexing agent.
28. The method of claim 26 where the electrophile is carbon dioxide.
29. The method of claim 26 where the electrophile is ethylbromoacetate.
- 25
30. A method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising
reacting a β -substituted pyrrole and a ketone to produce a β -substituted calix[n]pyrrole.
- 30
31. The method of claim 30 wherein the β -substituted calix[n]pyrrole is 3,4-dimethoxypyrrole, and the ketone is cyclohexanone, acetone, or pentan-3-one.
32. A method of making a β -substituted calix[n]pyrrole where n is 4, 5, 6, 7, or 8 comprising
producing a calix[n]pyrrole radical where n is 4, 5, 6, 7, or 8; and
35 reacting the radical with a radical acceptor molecule.
33. A method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, 8 comprising
reacting a pyrrole and a ketone in the presence of a Lewis acid.


34. The method of claim 33 wherein the Lewis acid is a high valent metal containing compound or boron trifluoride.
- 5 35. The method of claim 33 wherein n is 5.
36. The method of claim 33 wherein n is 6.
37. A method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising reacting a
10 pyrrole and a ketone in the presence of a heterogenous catalyst.
38. The method of claim 37 wherein the heterogenous catalyst is montmorillonite, zeolite, silica gel, alumina, or a cation exchange solid support.
- 15 39. The method of claim 37 wherein n is 5.
40. A method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising
reacting a pyrrole and a ketone in the presence of an anion template.
- 20 41. A method of making a calix[n]pyrrole wherein n is 4, 5, 6, 7, or 8 comprising
a) reacting a ketone-functionalized pyrrole and an alkyl metal to form a pyrrole alcohol,
b) condensing the pyrrole alcohol to form a cyclizable pyrrole oligomer; and
c) repeating step b) at least n-2 times to form a calix[n]pyrrole.
- 25 42. The method of claim 41 wherein the alkyl metal is a Grignard reagent or alkyl lithium.
43. The method of claim 41 wherein n is 5.
44. A method of making a solid-supported calix[n]pyrrole comprising
30 attaching a calix[n]pyrrole having a functionalized group to a solid support, the solid support reactive with the functionalized group, or to a tether-functionalized solid support, the tether reactive with the functionalized group.
45. A method of making a solid-supported calix[m]pyridino[n]pyrrole comprising
35 attaching a calix[m]pyridino[n]pyrrole having a functionalized group to a solid support, the solid support reactive with the functionalized group; or to a tether-functionalized solid support, the tether reactive with the functionalized group.

46. A method of making a solid-supported calix[m]pyridine comprising
attaching a calix[m]pyridine having a functionalized group to a solid support, the solid
support reactive with the functionalized group, or to a tether-functionalized solid support, the
tether reactive with the functionalized group.
- 5
47. A calix[n]pyrrole made by the method of claim 21, 23, 25, 26, 27, 30, 32, 33, 37, 40, 41, or
42.
48. A method of forming a complex of a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole and an
10 anion or a neutral molecule, comprising contacting the calix[n]pyrrole or the
calix[m]pyridino[n]pyrrole with the anion or neutral molecule under conditions effective to
allow the formation of the complex.
49. A method for separating a first molecule, a first anion, or first cation from a mixture of
15 molecules, anions or cations, comprising
obtaining a calix[n]pyrrole-, a calix[m]pyridino[n]pyrrole- or a calix[m]pyridine-
derivatized solid support; and
contacting the solid support with the mixture of molecules, anions or cations to separate
the first molecule, the first anion or the first cation.
- 20
50. The method of claim 49 wherein the solid support is in the form of a chromatography
column.
51. The method of claim 49 wherein the solid support is in the form of a capillary
25 electrophoresis tube.
52. The method of claim 49 wherein the contacting is in a batch process.
53. The method of claim 49 wherein the solid support is a calix[n]pyrrole- or a
30 calix[m]pyridino[n]pyrrole-solid support, the method is for separating a first anion, and the
first anion is fluoride, phosphate, a phosphorylated molecule, or nitrate.
54. The method of claim 49 wherein the solid support is a calix[n]pyrrole- or a
calix[m]pyridino[n]pyrrole-solid support, the method is for separating a first molecule, and the
35 first molecule is a nucleotide or an oligonucleotide.
55. The method of claim 49 wherein the method is for separating a first anion or a first cation,
and

the first anion is an atomic anion or an molecular anion, or
the first cation is a atomic cation or a molecular cation.

56. The method of claim 49 wherein the method is for separating a first molecule, and the first
5 molecule is a neutral, cationic or anionic aromatic molecule.

57. The method of claim 49 wherein the solid support is a calix[n]pyrrole solid support and the
calix[n]pyrrole has structure I



10

wherein

n is 4, 5, 6, 7 or 8;

when n is 4, p = q = r = s = 0, R₁ - R₁₆ are independently substituents as listed
in paragraph i) below, and R_A - R_D are independently substituents as
listed in paragraph ii) below;

15

when n is 5, p = 1, q = r = s = 0, R₁ to R₂₀ are independently substituents as
listed in paragraph i) below, and R_A - R_E are independently substituents
as listed in paragraph ii) below;

when n is 6, $p = q = 1$, $r = s = 0$, R_1 to R_{24} are independently substituents as listed in paragraph i) below, and $R_A - R_F$ are independently substituents as listed in paragraph ii) below;

when n is 7, $p = q = r = 1$, $s = 0$, R_1 to R_{28} are independently substituents as listed in paragraph i) below, and $R_A - R_G$ are independently substituents as listed in paragraph ii) below;

when n is 8, $p = q = r = s = 1$, R_1 to R_{32} are independently substituents as listed in paragraph i) below, and $R_A - R_H$ are independently substituents as listed in paragraph ii) below;

i) hydrogen, halide, hydroxyl, alkyl, alkenyl, alkynyl, aryl, alkylaryl, nitro, phospho, formyl, acyl, hydroxyalkyl, alkoxy, hydroxyalkoxy, hydroxyalkenyl, hydroxyalkynyl, saccharide, carboxy, carboxyalkyl, carboxyamide, carboxyamidealkyl, amino, amido, aminoalkyl, phosphoalkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, tetrahydropyran, tetrahydrothiapyran, thioalkyl, haloalkyl, haloalkenyl, haloalkynyl, alkyl ester, a site-directing molecule, a catalytic group, a reporter group, a binding agent, or a couple that is coupled to a site-directing molecule, to a catalytic group, to a reporter group, or to a binding agent;

ii) a lone pair of electrons, hydrogen, alkyl, aminoalkyl, alkylsulfone, carboxy alkyl, carboxyamidealkyl, phospho alkyl, alkyl sulfoxide, alkyl sulfone, alkyl sulfide, halo alkyl, aryl, N-oxide, dialkylamino, carbamate, or arylsulfonyl;

or

at least two substituents are coupled to form a bridged structure, and when coupled to form a bridged structure, nonbridged substituents are as defined herein in paragraph i) or ii) other than for bridged substituents;

wherein

at least one R substituent is attached to the solid support;

at least one pyrrole nitrogen is bonded to hydrogen; and

odd-numbered R-substituents are other than hydrogen.

58. The method of claim 49 wherein the method is for separating a first anion, the solid support is a calix[n]pyrrole- or a calix[m]pyridino[n]pyrrole-solid support and the first anion is pertechnetate.

59. The method of claim 49 wherein the method is for separating a first molecule, the solid support is a calix[n]pyrrole- or a calix[m]pyridino[n]pyrrole-solid support and the first molecule is a polyhalobiphenyl.
- 5 60. The method of claim 59 where the polyhalobiphenyl is a polychlorobiphenyl.
61. The method of claim 49 wherein the method is for separating a first anion, the solid support is a calix[n]pyrrole- or a calix[m]pyridino[n]pyrrole-solid support, and the first anion is a phosphate anion.
- 10 62. A method of transporting a molecular or anionic species through a membrane comprising incorporating a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole into the membrane; and contacting the membrane with the molecular or anionic species in the presence of a gradient of the molecular or anionic species or a counter gradient of a further species.
- 15 63. The method of claim 62 wherein the transporting results in the purification of the molecular or anionic species.
- 20 64. A method of binding a cation comprising contacting the cation with a calix[n]pyrrole or calix[m]pyridino[n]pyrrole having a cation-binding functionality, or with a calix[m]pyridine.
- 25 65. A method of removal of pertechnetate from pertechnetate-containing nuclear waste comprising contacting the waste with a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole to form a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole pertechnetate complex; and removing the complex from the water.
- 30 66. A method of removal of a polyhalobiphenyl compound from a polyhalobiphenyl containing environmental source comprising contacting the environmental source with a calix[n]pyrrole or a calix[m]pyridino[n]pyrrole to form a calix[n]pyrrole- or a calix[m]pyridino[n]pyrrole-polyhalobiphenyl complex; and removing the complex from the environmental source.
- 35 67. A chromatography column comprising a solid support bound to a calix[n]pyrrole where n is 4-8; to a calix[m]pyridino[n]pyrrole where $m+n=4-8$ and m is not 1 or 2; or to a calix[m]pyridine where m is 4-8.

68. A sensor comprising a solid support bound to a calix[n]pyrrole where n is 4-8; to a calix[m]pyridino[n]pyrrole where $m+n=4-8$ and m is not 1 or 2; or to a calix[m]pyridine where m is 4-8.
- 5
69. Use of a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine in the preparation of a pharmaceutical composition for use in *in vivo* or *ex vivo* treatment of body tissues.
- 10
70. The use of claim 69 wherein the treatment is *ex vivo* and is kidney dialysis.
71. The use of claim 69 wherein the treatment is removal of urea for treatment of gout.
72. An electropolymerizable calix[n]pyrrole, calix[m]pyridino[n]pyrrole, or calix[m]pyridine.
- 15
73. An anion-, cation-, or neutral molecule-selective electrode comprising
a conductive body,
a polymer, and
a calix[n]pyrrole, a calix[m]pyridino[n]pyrrole, or a calix[m]pyridine.
- 20
74. The electrode of claim 73 wherein the calix[n]pyrrole, the calix[m]pyridino[n]pyrrole, or the calix[m]pyridine is electropolymerized and forms the conductive body.
75. A method of electrochemical detection of an anion, a cation, or a neutral molecule
- 25 comprising
assembling the anion-, cation-, or neutral molecule-selective electrode of claim 73;
contacting the electrode with a solution of the anion, the cation, or the neutral molecule;
and
determining the presence or absence of the anion, the cation, or the neutral molecule.
- 30

100
ABSTRACT

5 The present invention provides calixpyrrole, calixpyridinopyrrole, and calixpyridine
macrocycles, having 4, 5, 6, 7, or 8 heterocyclic rings, as well as syntheses, derivatives,
conjugates, multimers, and solid supports thereof. Such macrocycles have proved to be
effective and selective ion- and neutral- molecule binding and ion- and neutral- molecule
separation agents. The macrocycles are fully *meso*-non-hydrogen-substituted porphyrinogens,
a few molecules of which were previously known but not recognized as possessing anion or
molecule binding properties. The binding mode is noncovalent, primarily that of hydrogen-
10 bonding, thereby providing a new mode for liquid chromatography, that of Hydrogen Bonding
Liquid Chromatography. Further useful applications of the macrocycles provided herein
include environmental remediation by removal of undesired ions or neutral molecules, and
removal of phosphate for kidney dialysis.




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4




FIG. 5

10/30




FIG. 6




FIG. 7




FIG. 8




FIG. 9

14/30






FIG.10






FIG. 12A




FIG. 12B




FIG. 13

18/30




FIG. 14




FIG. 15A




FIG. 15B




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20

25/30




FIG. 21




FIG. 22

27/30




FIG. 23




FIG. 24




FIG. 25

30/30




FIG. 26